arXiv:0710.1662v2 [hep-th] 15 Oct 2007

The Exact Renormalization Group[*]
– renormalization theory revisited –

Hidenori Sonoda[†]
Physics Department, Kobe University, Japan

September 2007

Abstract

We overview the entire renormalization theory, both perturbative and non-perturbative, by the method of the exact renormalization group (ERG). We emphasize particularly on the perturbative application of the ERG to the ϕ^4 theory and QED in the 4 dimensional euclidean space.

Contents

1 Introduction **3**

2 The Exact Renormalization Group (ERG) **4**
- 2.1 Notation 4
- 2.2 Derivation of the Polchinski ERG differential equation 6
- 2.3 Physical understanding of ERG 10
- 2.4 Perturbative renormalizability 11
- 2.5 Direct determination of the renormalized theory 13
- 2.6 Simple examples 14
- 2.7 The continuum limit in terms of a cutoff theory 18
- 2.8 ERG for the 1PI $\Gamma[\Phi]$ 19
- 2.9 Composite operators 22
- 2.10 Composite operators given in terms of the action 24
- 2.11 Beta functions and anomalous dimensions 29
- 2.12 Universality 31

3 Application to QED **34**
- 3.1 Perturbative construction 34
- 3.2 Ward identities 35
- 3.3 Perturbative construction 39

---

[*] Lectures given at the Feza Gürsey Institute, Istanbul, Turkey in 10 – 21 September, 2007.
[†] E-mail: `hsonoda@kobe-u.ac.jp`

3.4 BRST invariance 41
3.5 BRST with external sources 45
3.6 Explicit construction of $\mathcal{S}$ in terms of $\bar{S}$ 50

4 Application to YM theories 54
4.1 YM theories without sources 54
4.2 YM theories with sources 55

5 Non-perturbative aspects of the Wilson ERG 62
5.1 Generalized diffusion equation 62
5.2 Modification of the Polchinski ERG equation 68
5.3 Renormalization conditions 69
5.4 Rescaling of space 71
5.5 The Wilson-Fisher fixed point for $D = 3$ 74
5.6 Perturbative determination of critical exponents 75

1 Introduction

The subject of the following series of lectures is the exact renormalization group (ERG), also known as the Wilsonian renormalization group. Ken Wilson invented it in the early 70's to understand the physics behind renormalization of quantum field theory. Quantum field theory had seen phenomenal success in its applications to QED via renormalization. But many, if not most, people felt uneasy about renormalization. It was quite common to regard renormalization as a clever mathematical trick to hide what is not understood under the rug. Not many had even imagined there was physics behind the procedure of renormalization.

To understand the physics of renormalization, Wilson introduced RG flows in the space of theories. Using such important notions as criticality, fixed points, and relevant and marginal parameters, the continuum limit is defined as the RG trajectories flowing out of a UV fixed point. The continuum limit can be constructed by using an almost critical theory by using the scale dimensions of relevant parameters. All this has been explained in the classic review article of Wilson and Kogut [1], especially in sect. 12.

In the following lectures, our emphasis is on the perturbative applications of ERG. Only in the last part, we discuss the non-perturbative aspects of ERG. Even there the emphasis is on the relation of Wilson's original ERG differential equation to Polchinski's which can be applied much more easily to perturbation theory.

The lectures are organized into four parts. In the first part we review the application of ERG to the most relevant aspects of perturbative renormalization theory using the ϕ^4 theory for concreteness: We derive Polchinski's version of the ERG differential equation that describes the cutoff dependence of the Wilson action.[2] We then show how to incorporate perturbative renormalizability as asymptotic behaviors of the action as the cutoff is raised toward infinity. We show how to determine the cutoff action by solving the ERG differential equation under the asymptotic conditions. Using composite operators as an essential tool, we use ERG to derive the RG equations of the parameters of the theory, and to show universality.

In the second and third parts, we show how a momentum cutoff can be consistent with gauge symmetry. In the second part we take a bottom-to-up approach to construct a cutoff action of QED: we first construct an action that reproduces the Ward identities of the correlation functions, and then introduce Faddeev-Popov ghosts to introduce a BRST invariant action. We finally introduce sources that generate BRST transformations to build a fully BRST invariant formalism with nilpotent BRST transformations. The discussions of YM theories in the third part are relatively brief. We outline a construction of a fully BRST invariant cutoff action with sources. The sources are necessary to show the possibility of satisfying BRST invariance. Once the possibility is known, practical calculations can be done without the sources. Part of the new results contained in the second and third parts of the lectures have been obtained in collaboration with Y. Igarashi and K. Itoh.

Finally, in the last part we discuss the non-perturbative aspects of ERG. In the first subsection we give a non-perturbative derivation of a general class of ERG differential equations. We then relate Wilson's original ERG differential equation to Polchinski's explicitly via an integral formula. The two sets of ERG differential equations are completely equivalent. We then show how to modify Polchinski's ERG differential equation to accommodate anomalous dimensions. We end the last part by going back to perturbation theory, computing the critical exponents of the Wilson-Fisher fixed point using ERG.

Best efforts have been made to make the lectures as self-contained as possible. The reader is warned, though, that my style may be somewhat idiosyncratic; the ERG is a rich field, and it can accommodate widely varying viewpoints. The aim of the following lectures is to give a coherent overview of the applications of ERG to renormalization theory, but not to give a balanced review of the field of ERG. For the latter and also for further references, consultation with the existing reviews [3, 4, 5, 6, 7, 8, 9, 10, 11, 12] is strongly recommended. The other references I cite are kept minimal except for the inclusion of refs.[17, 18] which happen to be relevant to my current research interest.

2 The Exact Renormalization Group (ERG)

2.1 Notation

We consider a real scalar theory in D-dimensional euclidean space. The scalar field is given by

$$\phi(r) = \int_p \phi(p)\, \mathrm{e}^{ipr} \qquad \left(\int_p \equiv \int \frac{d^D p}{(2\pi)^D} \right)$$

where

$$pr \equiv \sum_{\mu=1}^{D} p_\mu r_\mu$$

Given an action $S[\phi]$, we define the correlation functions by

$$\langle \phi(p_1) \cdots \phi(p_n) \rangle_S \cdot (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_n) \equiv \frac{\int [d\phi]\, \phi(p_1) \cdots \phi(p_n)\, \mathrm{e}^{S[\phi]}}{\int [d\phi]\, \mathrm{e}^{S[\phi]}}$$

Note that the weight of the functional integral is e^S, not e^{-S}. For the higher-point functions $n > 2$, we usually consider only the connected part:

$$\langle \phi(p_1) \cdots \phi(p_n) \rangle \overset{n>2}{\Longrightarrow} \langle \phi(p_1) \cdots \phi(p_n) \rangle^{\mathrm{connected}}$$

We consider a real scalar theory whose propagator is given by

$$\boxed{\frac{K(p/\Lambda)}{p^2 + m^2}}$$

where $K(p)$ is a positive decreasing function of p^2 with the property that

$$K(p)\left\{\begin{array}{ll} = 1 & (p^2 < 1) \\ \to 0 & (p^2 \to \infty) \end{array}\right.$$



Figure 1: The cutoff function K and its derivative Δ

The propagation of the momentum modes with $p^2 > \Lambda^2$ is suppressed sufficiently strongly so that the loop integrals for each Feynman diagram (to be introduced shortly) become well defined, i.e., **free of UV divergences**. For later conveniences, we also define the logarithmic derivative

$$\Delta(p) \equiv -2p^2 \frac{dK(p^2)}{dp^2}$$

which vanishes for $p^2 < 1$, and is positive for $p^2 > 1$ with a peak near $p^2 = 1$.

For example, we can take

$$K(p) \equiv \left\{\begin{array}{cl} 1 & (p^2 < 1) \\ 1 - \mathrm{e}^{-\frac{1}{(p^2-1)^n}} & (p^2 > 1) \end{array}\right.$$

so that

$$\Delta(p) = \left\{\begin{array}{cl} 0 & (p^2 < 1) \\ \frac{2np^2}{(p^2-1)^{n+1}}\, \mathrm{e}^{-\frac{1}{(p^2-1)^n}} & (p^2 > 1) \end{array}\right.$$

Here, n is a big enough integer. For $D = 4$, the worst divergence is quadratic, so $n = 2$ suffices.

Let us suppose that the action is given by

$$S[\phi] = S_{\mathrm{free}}[\phi] + S_{\mathrm{int}}[\phi]$$

where the free action

$$S_{\mathrm{free}}[\phi] \equiv -\frac{1}{2} \int_p \phi(p)\phi(-p) \frac{p^2 + m^2}{K(p/\Lambda)}$$

defines the propagator given above, and the interaction part is given by

$$S_{\mathrm{int}}[\phi] \equiv \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1,\cdots,p_{2n}} \phi(p_1)\cdots\phi(p_{2n})\, \mathcal{V}_{2n}(p_1,\cdots,p_{2n}) \cdot (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_{2n})$$

We have assumed the $\mathbf{Z}_2$ invariance for simplicity.[1] We assume that the vertex $\mathcal{V}_{2n}$ is local: For our purposes we only need to assume that $\mathcal{V}_{2n}$ can be expanded in powers of small momenta. We denote the $2n$-point vertex graphically by a blob: The perturbative expansion of a correlation function is given by the sum over Feynman diagrams with propagators and vertices.



Figure 2: A blob represents a $2n$-point vertex $\mathcal{V}_{2n}$.

For example, the diagrams in Fig. 3 contribute to the two-point function: and those in Fig. 4 contribute to the four-point function.



Figure 3: These contribute to the two-point function.



Figure 4: These contribute to the four-point function.

2.2 Derivation of the Polchinski ERG differential equation

We solve the following problem: we decrease the momentum cutoff Λ infinitesimally to $\Lambda \mathrm{e}^{-\Delta t}$.

$$\boxed{\Lambda \longrightarrow \Lambda \mathrm{e}^{-\Delta t}}$$

We wish to compensate the change by appropriate changes in the vertices so that the new propagator and vertices give the same correlation functions.

$$\boxed{\begin{array}{c}\Lambda\\ \mathcal{V}_{2n}\end{array}} \quad \overset{\text{equivalent}}{\Leftrightarrow} \quad \boxed{\begin{array}{c}\Lambda \mathrm{e}^{-\Delta t}\\ \mathcal{V}'_{2n}\end{array}}$$

---

[1] We also assume that the symmetry is not broken spontaneously. Hence, we will only consider even-point correlation functions.

To solve this problem, we consider an arbitrary Feynman diagram, and classify the propagators into the three types:[2]

1. those connecting two different vertices (type A)
2. those connecting a single vertex (type B)
3. those attaching an external line to a vertex (type C)



Figure 5: The three types of propagators

The original propagator is related to the new propagator as follows:

$$\frac{K(p/\Lambda)}{p^2+m^2} = \frac{K(p/(\Lambda \mathrm{e}^{-\Delta t}))}{p^2+m^2} + \Delta t\, \frac{\Delta(p/\Lambda)}{p^2+m^2}$$

We consider a Feynman graph with the original propagators and vertices. Then, we replace each propagator by the sum on the right-hand side. At first order in Δt, we get a collection of Feynman diagrams in which one of the propagators is replaced by

$$\Delta t\, \frac{\Delta(p/\Lambda)}{p^2+m^2}$$

If the propagator is either type A or B, we can interpret it as a correction to the relevant vertex. (Fig. 6) Hence, the type A and B propagators modify the $2n$-point vertex to

$$\begin{aligned} &\mathcal{V}_{2n}(p_1,\cdots,p_{2n}) \Longrightarrow \mathcal{V}'_{2n}(p_1,\cdots,p_{2n}) \equiv \mathcal{V}_{2n}(p_1,\cdots,p_{2n}) \\ &\quad +\Delta t \sum_{\text{partitions}} \mathcal{V}_{2k}(p_{i_1},\cdots,p_{i_{2k-1}},q)\, \underbrace{\frac{\Delta(q/\Lambda)}{q^2+m^2}}_{\text{type A}}\, \mathcal{V}_{2(n+1-k)}(-q,p_{i_{2k}},\cdots,p_{i_{2n}}) \\ &\quad +\Delta t\, \frac{1}{2} \int_q \underbrace{\frac{\Delta(q/\Lambda)}{q^2+m^2}}_{\text{type B}}\, \mathcal{V}_{2(n+1)}(q,-q,p_1,\cdots,p_{2n}) \end{aligned}$$

where the momentum q of the type A propagator is fixed as

$$q \equiv -\left(p_{i_1}+\cdots+p_{i_{2k-1}}\right) = p_{i_{2k}}+\cdots+p_{i_{2n}}$$

but the momentum q of the type B propagator must be integrated over. Since

---

[2] There is actually one exception to this classification: the free propagator contributing to the two-point function. We will take care of this later.



Figure 6: Type A and B propagators modify the vertex $\mathcal{V}_{2n}$.

$\Delta(q/\Lambda)$ is non-vanishing only for $q^2 > \Lambda^2$, we can consider the change of the vertices as local, i.e., in coordinate space the change takes place over a distance of order $1/\Lambda$.

The change in the type C propagators can be corrected simply by the multiplicative factor

$$\frac{K(p_i/\Lambda)}{K(p_i/(\Lambda \mathrm{e}^{-\Delta t}))} = 1 + \Delta t \, \frac{\Delta(p_i/\Lambda)}{K(p_i/\Lambda)}$$

for each external line.

Using the modified vertices, we define a new action by

$$\begin{aligned} S'[\phi] &= S[\phi] + \delta S[\phi] \\ &\equiv -\frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2+m^2}{K(p/(\Lambda \mathrm{e}^{-\Delta t}))} \\ &\quad + \sum_{n=1}^{\infty} \frac{1}{(2n)!}\int_{p_1,\cdots,p_{2n}} \mathcal{V}'_{2n}(p_1,\cdots,p_{2n}) \cdot (2\pi)^D \delta^{(D)}(p_1+\cdots+p_{2n}) \end{aligned}$$

The diagrammatic analysis above implies

$$\langle \phi(p_1)\cdots\phi(p_{2n})\rangle_S = \prod_{i=1}^{2n} \frac{K(p_i/\Lambda)}{K(p_i/(\Lambda \mathrm{e}^{-\Delta t}))} \cdot \langle \phi(p_1)\cdots\phi(p_{2n})\rangle_{S'}$$

Hence,

$$\langle \phi(p_1)\cdots\phi(p_{2n})\rangle_S - \langle \phi(p_1)\cdots\phi(p_{2n})\rangle_{S+\delta S} = \Delta t \sum_{i=1}^{2n} \frac{\Delta(p_i/\Lambda)}{K(p_i/\Lambda)} \cdot \langle \phi(p_1)\cdots\phi(p_{2n})\rangle_S$$

In fact the above relation is valid only for $2n \geq 4$. For the two-point function, the first term in its perturbative expansion is the free propagator. It cannot be classified into any of the three types.

$$\langle \phi(p)\phi(-p)\rangle_S = \qquad \text{———} \qquad + \text{diagrams with at least one vertex}$$

Hence, it is the difference

$$\langle \phi(p)\phi(-p)\rangle_S - \frac{K(p/\Lambda)}{p^2+m^2}$$

which contains only the propagators of the three types. Therefore, we obtain

$$\langle\phi(p)\phi(-p)\rangle_S - \frac{K(p/\Lambda)}{p^2+m^2} = \frac{K(p/\Lambda)^2}{K(p/(\Lambda\mathrm{e}^{-\Delta t}))^2} \cdot \left(\langle\phi(p)\phi(-p)\rangle_{S'} - \frac{K(p/(\Lambda\mathrm{e}^{-\Delta t}))}{p^2+m^2} \right)$$

We can rewrite this as

$$\begin{aligned} &\langle\phi(p)\phi(-p)\rangle_S - \langle\phi(p)\phi(-p)\rangle_{S+\delta S} \\ &= \Delta t \left[2\frac{\Delta(p/\Lambda)}{K(p/\Lambda)} \cdot \langle\phi(p)\phi(-p)\rangle_S - \frac{\Delta(p/\Lambda)}{K(p/\Lambda)} \frac{K(p/\Lambda)}{p^2+m^2} \right] \end{aligned}$$

The expression of the new action $S' = S + \delta S$, which is equivalent to the original action S, is easy to understand graphically, but quite complicated to write down as a formula. By using a functional notation, δS_{int} (the interaction part of δS) can be expressed quite simply:

$$\delta S_{\mathrm{int}} = \Delta t \int_q \frac{\Delta(q/\Lambda)}{q^2+m^2} \frac{1}{2} \left\{ \frac{\delta S_{\mathrm{int}}}{\delta\phi(q)} \frac{\delta S_{\mathrm{int}}}{\delta\phi(-q)} + \frac{\delta^2 S_{\mathrm{int}}}{\delta\phi(q)\delta\phi(-q)} \right\}$$

HW#1: Convince yourself this is the same as the equation we have obtained using Feynman graphs.

Writing the original S as $S(\Lambda)$ and S' as $S(\Lambda\mathrm{e}^{-\Delta t})$, we obtain

$$\delta S_{\mathrm{int}} = \Delta t \cdot \left(-\Lambda \frac{\partial}{\partial\Lambda} \right) S_{\mathrm{int}}$$

Hence, $S_{\mathrm{int}}(\Lambda)$ satisfies the following differential equation:

$$\boxed{-\Lambda \frac{\partial S_{\mathrm{int}}(\Lambda)}{\partial\Lambda} = \int_q \frac{\Delta(q/\Lambda)}{q^2+m^2} \frac{1}{2} \left\{ \frac{\delta S_{\mathrm{int}}}{\delta\phi(q)} \frac{\delta S_{\mathrm{int}}}{\delta\phi(-q)} + \frac{\delta^2 S_{\mathrm{int}}}{\delta\phi(q)\delta\phi(-q)} \right\}}$$

This was first derived by Polchinksi.[2] The transformation from $S(\Lambda)$ to $S(\Lambda\mathrm{e}^{-t})$, where $t > 0$, is called an **exact renormalization group (ERG)** transformation: it is exact since we have not introduced any approximation.

Using

$$\begin{aligned} \delta S_{\mathrm{free}} &= S_{\mathrm{free}}(\Lambda\mathrm{e}^{-\Delta t}) - S_{\mathrm{free}}(\Lambda) \\ &= -\frac{1}{2} \int_p \phi(p)\phi(-p) \left(p^2+m^2\right) \left(\frac{1}{K(p/(\Lambda\mathrm{e}^{-\Delta t}))} - \frac{1}{K(p/\Lambda)} \right) \\ &= -\Delta t \frac{1}{2} \int_p \phi(p)\phi(-p) \left(p^2+m^2\right) \frac{\Delta(p/\Lambda)}{K(p/\Lambda)^2} \end{aligned}$$

further, we can rewrite the Polchinski differential equation for the entire action as follows:

$$\boxed{\begin{aligned} -\Lambda\frac{\partial}{\partial\Lambda} S(\Lambda) = \int_q \frac{\Delta(q/\Lambda)}{q^2+m^2} \Bigg[& \frac{q^2+m^2}{K(q/\Lambda)} \phi(q) \frac{\delta S(\Lambda)}{\delta\phi(q)} \\ & + \frac{1}{2} \left\{ \frac{\delta S(\Lambda)}{\delta\phi(q)} \frac{\delta S(\Lambda)}{\delta\phi(-q)} + \frac{\delta^2 S(\Lambda)}{\delta\phi(q)\delta\phi(-q)} \right\} \Bigg] \end{aligned}}$$

HW#2: Derive this from Polchinski's. Alternatively, derive Polchinski's from this. (You need to ignore an infinite constant independent of ϕ. An additive constant to S does not affect physics.)

To summarize, the correlation functions satisfy the following ERG differential equations:

$$\begin{aligned}
-\Lambda\frac{\partial}{\partial\Lambda}\left\langle\phi(p)\phi(-p)\right\rangle_S &= \left\langle\phi(p)\phi(-p)\left(-\Lambda\frac{\partial S}{\partial\Lambda}\right)\right\rangle_S^{\text{connected}}\\
&= -2\frac{\Delta(p/\Lambda)}{K(p/\Lambda)}\left\langle\phi(p)\phi(-p)\right\rangle_S + \frac{\Delta(p/\Lambda)}{p^2+m^2}\\
-\Lambda\frac{\partial}{\partial\Lambda}\left\langle\phi(p_1)\cdots\phi(p_{2n})\right\rangle &= \left\langle\phi(p_1)\cdots\phi(p_{2n})\left(-\Lambda\frac{\partial S}{\partial\Lambda}\right)\right\rangle_S^{\text{connected}}\\
&= -\sum_{i=1}^{2n}\frac{\Delta(p_i/\Lambda)}{K(p_i/\Lambda)}\cdot\left\langle\phi(p_1)\cdots\phi(p_{2n})\right\rangle_S
\end{aligned}$$

2.3 Physical understanding of ERG

To gain physical (or intuitive) understanding of ERG, let us forget about UV divergences or any other difficulties our theory might face with, for the time being.

The standard propagator can be expressed as the sum:

$$\frac{1}{p^2+m^2} = \frac{K(p/\Lambda)}{p^2+m^2} + \frac{1-K(p/\Lambda)}{p^2+m^2}$$

The first term on the right is the cutoff propagator for the low momentum modes. The second term is the propagator for the high momentum modes.

Given a Feynman diagram with the standard propagators and elementary vertices, we substitute the above sum into each propagator. The substitution generates multiple diagrams in which some propagators are the low-momentum propagators, and the rest are high-momentum propagators. Then, we pay attention to those parts of the Feynman diagram connected only by high-momentum propagators. The entire diagram has these connected parts connected further by low-momentum propagators. (Fig. 7) Suppose a connected part has $2n$ ex-



Figure 7: Each blob contains only high-momentum propagators inside.



Figure 8: $\mathcal{V}_{2n}(\Lambda)$ consists of graphs whose internal lines carry only high momenta above Λ.

ternal lines. Then, the connected part can be interpreted as a contribution to the vertices $\mathcal{V}_{2n}(\Lambda)$ of the cutoff theory $S(\Lambda)$. (Fig. 8) Thus, we put all the short-distance physics (shorter than the length $\frac{1}{\Lambda}$) into the vertices.

Now, the problem is the potential UV divergence of the diagrams with the high-momentum propagators. For example, the 1st order contribution to $\mathcal{V}_2$ is given by



$$\stackrel{?}{=} -\frac{\lambda}{2}\int_q \frac{1-K(q/\Lambda)}{q^2+m^2}$$

which is quadratically divergent since $K(q/\Lambda) \to 0$ as $q^2 \to \infty$. We will define the diagram, not by the naive loop integral, but as a finite solution to the ERG differential equation. It is the role of ERG to make sense out of the naively UV divergent Feynman diagrams.

2.4 Perturbative renormalizability

Polchinski introduced his differential equation in order to simplify the proof of perturbative renormalizability of the ϕ^4 theory in $D = 4$. We will not follow his proof, but merely sketch his ideas here. We start from a bare action defined at a large momentum scale Λ:

$$\begin{aligned} S_\Lambda(\Lambda) = &-\frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2+m^2}{K(p/\Lambda)} \\ &-\frac{1}{2}\int_p \phi(p)\phi(-p)\left(\Delta z_0 \cdot p^2 + \Delta m_0^2\right) - \frac{\lambda_0}{4!}\int_{p_1+p_2+p_3+p_4=0} \phi(p_1)\cdots\phi(p_4) \end{aligned}$$

We then reduce the cutoff Λ to an arbitrary but fixed finite momentum scale μ to obtain an equivalent action $S_\Lambda(\mu)$. Here we have added a suffix Λ to remind ourselves that we started from the scale Λ. We wish to introduce appropriate Λ dependence to the bare parameters

$$\Delta z_0(\Lambda), \quad \Delta m_0^2(\Lambda), \quad \lambda_0(\Lambda)$$



Figure 9: Renormalizability amounts to the existence of $\lim_{\Lambda\to\infty} S_\Lambda(\mu)$.

such that the continuum limit

$$\bar{S}(\mu) = \lim_{\Lambda\to\infty} S_\Lambda(\mu)$$

exists. (Fig. 9) This is **perturbative** renormalizability since we prove this, order by order in a coupling constant λ. Polchinski has proven this using his differential equation.

Why is this renormalizability? ERG guarantees that the correlation functions we compute with $\bar{S}(\mu)$ is the same as those with the bare action $S_\Lambda(\Lambda)$. In the limit $\Lambda \to \infty$, the theory is defined all the way up to the zero distance scale, and the correlation functions are independent of Λ. This is what we usually call renormalizability.

Typically, the bare parameters have the following Λ dependence:

$$\begin{cases} \Delta m_0^2(\Lambda) &= \Lambda^2 \cdot g\left(\frac{m^2}{\mu^2}, \ln\frac{\Lambda}{\mu}, \lambda\right) + m^2 \cdot z_m\left(\frac{m^2}{\mu^2}, \ln\frac{\Lambda}{\mu}, \lambda\right) \\ \Delta z_0(\Lambda) &= z\left(\frac{m^2}{\mu^2}, \ln\frac{\Lambda}{\mu}, \lambda\right) \\ \lambda_0(\Lambda) &= z_\lambda\left(\frac{m^2}{\mu^2}, \ln\frac{\Lambda}{\mu}, \lambda\right) \end{cases}$$

At each order of λ, the coefficient functions are finite degree polynomials of $\ln\frac{\Lambda}{\mu}$. The so-called **mass independent** scheme is distinguished by the absence of the dependence on $\frac{m^2}{\mu^2}$ in the coefficient functions. Hence, in a mass independent scheme, we find

$$\begin{cases} \Delta m_0^2(\Lambda) &= \Lambda^2 \cdot g\left(\ln\frac{\Lambda}{\mu}, \lambda\right) + m^2 \cdot z_m\left(\ln\frac{\Lambda}{\mu}, \lambda\right) \\ \Delta z_0(\Lambda) &= z\left(\ln\frac{\Lambda}{\mu}, \lambda\right) \\ \lambda_0(\Lambda) &= z_\lambda\left(\ln\frac{\Lambda}{\mu}, \lambda\right) \end{cases}$$

2.5 Direct determination of the renormalized theory

We now make a crucial observation. The action $\bar{S}(\mu)$, which defines the renormalized theory or the continuum limit, satisfies the ERG differential equation of Polchinski with respect to μ:

$$\boxed{\begin{aligned} -\mu \frac{\partial \bar{S}(\mu)}{\partial \mu} = & \int_q \frac{\Delta(q/\mu)}{q^2+m^2} \left[\frac{q^2+m^2}{K(q/\mu)} \phi(q) \frac{\delta \bar{S}}{\delta \phi(q)} \right. \\ & \left. + \frac{1}{2} \left\{ \frac{\delta \bar{S}}{\delta \phi(q)} \frac{\delta \bar{S}}{\delta \phi(-q)} + \frac{\delta^2 \bar{S}}{\delta \phi(q) \delta \phi(-q)} \right\} \right] \end{aligned}}$$

Hence, we should be able to determine $\bar{S}(\mu)$ directly without starting from $S_\Lambda(\Lambda)$ by solving the differential equation. But as is the case with any differential equation, we need what amounts to an initial condition. We cannot use $\bar{S}(\Lambda) = S_\Lambda(\Lambda)$, because $S_\Lambda(\Lambda)$ will not lie on the ERG trajectory of $\bar{S}(\Lambda)$. Instead we adopt the following:

1. $\mathcal{V}_{2n}(\Lambda; p_1, \cdots, p_{2n})$ vanishes as $\Lambda \to \infty$ for $2n \geq 6$
2. $\mathcal{V}_2$ satisfies the asymptotic condition

   $$\begin{aligned} \mathcal{V}_2(\Lambda; p, -p) \quad \overset{\Lambda\to\infty}{\longrightarrow} \quad & \Lambda^2 a_2(m^2/\mu^2, \ln \Lambda/\mu; \lambda) \\ & + m^2 b_2(m^2/\mu^2, \ln \Lambda/\mu; \lambda) + p^2 c_2(m^2/\mu^2, \ln \Lambda/\mu; \lambda) \end{aligned}$$

   and the conditions at $\Lambda = \mu$, where μ is fixed but arbitrary:

   $$\left\{ \begin{array}{c} \mathcal{V}_2(\mu; 0, 0) = 0 \\ \frac{\partial}{\partial p^2} \mathcal{V}_2(\mu, p, -p)\Big|_{p^2=0} = 0 \end{array} \right.$$

3. $\mathcal{V}_4$ satisfies the asymptotic condition

   $$\mathcal{V}_4(\Lambda; p_1, \cdots, p_4) \overset{\Lambda\to\infty}{\longrightarrow} a_4(m^2/\mu^2, \ln \Lambda/\mu; \lambda)$$

   and the condition at $\Lambda = \mu$:

   $$\mathcal{V}_4(\mu; 0, \cdots, 0) = -\lambda$$

The asymptotic conditions are crucial for renormalizability. The two conditions on $\mathcal{V}_2$ have to do with mass and wave function renormalization, and the condition on $\mathcal{V}_4$ normalizes the coupling λ.

The above scheme is not mass independent. For mass independence, we can alternatively adopt the following conditions:

1. $\mathcal{V}_{2n}(\Lambda; p_1, \cdots, p_{2n})$ vanishes as $\Lambda \to \infty$ for $2n \geq 6$

2. $\mathcal{V}_2$ satisfies the asymptotic condition

$$\mathcal{V}_2(\Lambda;p,-p) \overset{\Lambda\to\infty}{\longrightarrow} \Lambda^2 a_2(\ln\Lambda/\mu;\lambda) + m^2 b_2(\ln\Lambda/\mu;\lambda) + p^2 c_2(\ln\Lambda/\mu;\lambda)$$

and the conditions at $\Lambda=\mu$:

$$\begin{aligned} \frac{\partial}{\partial m^2}\mathcal{V}_2(\mu;0,0)\Big|_{m^2=0} &= 0 \\ \frac{\partial}{\partial p^2}\mathcal{V}_2(\mu;p,-p)\Big|_{m^2=p^2=0} &= 0 \end{aligned}$$

which implies

$$b_2(0;\lambda) = c_2(0;\lambda) = 0$$

3. $\mathcal{V}_4$ satisfies the asymptotic condition

$$\mathcal{V}_4(\Lambda;p_1,\cdots,p_4) \overset{\Lambda\to\infty}{\longrightarrow} a_4(\ln\Lambda/\mu;\lambda)$$

and the condition at $\Lambda=\mu$:

$$\mathcal{V}_4(\mu;0,\cdots,0)\Big|_{m^2=0} = -\lambda$$

which implies

$$a_4(0;\lambda) = 0$$

Note the absence of m^2 dependence in the coefficient functions a_2, b_2, c_2, and a_4.

Hence, to show renormalizability of ϕ^4 theory in $D=4$, we must show that given m^2, λ, and an arbitrary scale μ, the solution $\bar{S}(\Lambda)$ of the Polchinski differential equation is determined uniquely. This was done for the mass independent scheme in [14].[3]

2.6 Simple examples

Let us compute the vertices for $D=4$ to see how the conditions introduced above determine the vertices uniquely.

First order in λ

The four-point vertex is given by

$$\mathcal{V}_4^{(0)}(p_1,\cdots,p_4) = -\lambda$$

This is the starting point of perturbative calculations.

The two-point vertex $\mathcal{V}_2^{(1)}(\Lambda)$ satisfies the differential equation

$$-\Lambda\frac{\partial\mathcal{V}_2^{(1)}(\Lambda)}{\partial\Lambda} = \frac{1}{2}\int_q \frac{\Delta(q/\Lambda)}{q^2+m^2}\mathcal{V}_4^{(0)}(q,-q,p,-p) = -\lambda\frac{1}{2}\int_q \frac{\Delta(q/\Lambda)}{q^2+m^2}$$

This can be expressed diagrammatically as

---

[3] From now on, we only consider $\bar{S}$, and omit the bar.



where we use the notation

 $$= \frac{\Delta(q/\Lambda)}{q^2+m^2}$$

We can compute the right-hand side further as

$$\begin{aligned} &= -\lambda\frac{1}{2}\int_q \Delta(q/\Lambda)\left(\frac{1}{q^2}-\frac{m^2}{q^4}+\frac{m^4}{q^4(q^2+m^2)}\right)\\ &= -\lambda\frac{1}{2}\left(\Lambda^2\int_q\frac{\Delta(q)}{q^2}-m^2\int_q\frac{\Delta(q)}{q^4}+m^4\int_q\Delta(q/\Lambda)\frac{1}{q^4(q^2+m^2)}\right)\end{aligned}$$

Integrating this over Λ, we obtain

$\mathcal{V}_2^{(1)}(\Lambda) =$ 

$$\begin{aligned} = \; -\frac{\lambda}{2}\Bigg(&-\frac{\Lambda^2}{2}\int_q\frac{\Delta(q)}{q^2}+m^2\ln\frac{\Lambda}{\mu}\int_q\frac{\Delta(q)}{q^4}+m^4\int_q\frac{1-K(q/\Lambda)}{q^4(q^2+m^2)}\\ &+\frac{\mu^2}{2}\int_q\frac{\Delta(q)}{q^2}-m^4\int_q\frac{1-K(q/\mu)}{q^4(q^2+m^2)}\Bigg)\end{aligned}$$

which satisfies the condition

$$\mathcal{V}_2^{(1)}(\Lambda=\mu)=0$$

Alternatively, in the mass independent scheme, we obtain

$$\mathcal{V}_2^{(1)}(\Lambda) = -\frac{\lambda}{2}\left(-\frac{\Lambda^2}{2}\int_q\frac{\Delta(q)}{q^2}+m^2\ln\frac{\Lambda}{\mu}\int_q\frac{\Delta(q)}{q^4}+m^4\int_q\frac{1-K(q/\Lambda)}{q^4(q^2+m^2)}\right)$$

which has the simple asymptotic behavior

$$\mathcal{V}_2^{(1)}(\Lambda)\to -\frac{\lambda}{2}\left(-\frac{\Lambda^2}{2}\int_q\frac{\Delta(q)}{q^2}+m^2\ln\frac{\Lambda}{\mu}\int_q\frac{\Delta(q)}{q^4}\right)$$

Second order in λ

The ERG differential equation for the six-point vertex is given diagrammatically as

$$-\Lambda\frac{\partial}{\partial\Lambda}\mathcal{V}_6^{(0)}(\Lambda;p_1,\cdots,p_6) =$$  $+\,2$ permutations

Solving this, we obtain

$$\mathcal{V}_6^{(0)}(\Lambda;p_1,\cdots,p_6) =$$



$$+\,2 \text{ permutations}$$

$$= \lambda^2 \left(\frac{1-K((p_1+p_2+p_3)/\Lambda)}{(p_1+p_2+p_3)^2+m^2} + 2 \text{ permutations} \right)$$

This vanishes as $\Lambda \to \infty$.

At one-loop the four-point vertex satisfies the ERG differential equation:

$$-\Lambda \frac{\partial}{\partial \Lambda} \mathcal{V}_4^{(1)}(\Lambda;p_1,\cdots,p_4) = \sum_{i=1}^{4} \Bigg($$



$$+$$



$$\Bigg)$$

$$+$$



$$+2 \text{ permutations}$$

The first part on the right-hand side can be integrated to

$$(-\lambda) \sum_{i=1}^{4} \frac{1-K(p_i/\Lambda)}{p_i^2+m^2} \cdot \mathcal{V}_2^{(1)}(\Lambda)$$

The first graph on the second part gives

$$\lambda^2 \int_q \frac{1-K((q+p_1+p_2)/\Lambda)}{(q+p_1+p_2)^2+m^2} \frac{\Delta(q/\Lambda)}{q^2+m^2} = \lambda^2 \int_q \left[\frac{(1-K(q/\Lambda))\,\Delta(q/\Lambda)}{q^4} \right.$$
$$\left. + \frac{1-K((q+p_1+p_2)/\Lambda)}{(q+p_1+p_2)^2+m^2} \frac{\Delta(q/\Lambda)}{q^2+m^2} - \frac{(1-K(q/\Lambda))\,\Delta(q/\Lambda)}{q^4} \right]$$

The first integral on the right gives a constant

$$\int_q \frac{\Delta(q)\,(1-K(q))}{q^4}$$

The remaining integral can be integrated over Λ to give a finite result:

$$\frac{1}{2} \int_q \left[\frac{1-K((q+p_1+p_2)/\Lambda)}{(q+p_1+p_2)^2+m^2} \frac{1-K(q/\Lambda)}{q^2+m^2} - \frac{(1-K(q/\Lambda))^2}{q^4} \right]$$

Hence, we obtain



$$= -\lambda^2 \ln \frac{\Lambda}{\mu} \int_q \frac{\Delta(q)\,(1-K(q))}{q^4}$$
$$+ \frac{\lambda^2}{2} \int_q \left[\frac{1-K((q+p_1+p_2)/\Lambda)}{(q+p_1+p_2)^2+m^2} \frac{1-K(q/\Lambda)}{q^2+m^2} - \frac{(1-K(q/\Lambda))^2}{q^4} \right]$$

where we have adopted the mass independence. This has the asymptotic property



$$\overset{\Lambda\to\infty}{\longrightarrow} -\lambda^2 \ln\frac{\Lambda}{\mu}\int_q \frac{\Delta(q)\,(1-K(q))}{q^4}$$

HW#3: Evaluate the one-loop four-point vertex using the condition

$$\mathcal{V}_4(\mu;0,\cdots,0) = -\lambda$$

instead of the mass independence. You will see that the asymptotic behavior acquires m^2 dependence.

Integrals

In general the integrals involving the cutoff function K depend on the choice of K. Certain integrals are universal, however. Let us first consider

$$I(n) \equiv \int_q \Delta(q)\frac{K(q)^n}{q^4}$$

in $D = 4$. This is integrable, since the integrand vanishes exponentially as $q^2 \to \infty$, and also vanishes for $0 < q^2 < 1$. Since

$$\Delta(q) = -2q^2\frac{dK(q)}{dq^2}$$

we obtain

$$\Delta(q)K(q)^n = \frac{1}{n+1}(-2q^2)\frac{dK(q)^{n+1}}{dq^2}$$

Hence,

$$\begin{aligned} I_n &= \frac{2\pi^2}{(2\pi)^4}\frac{1}{n+1}\int_0^\infty \frac{q^2dq^2}{2}\frac{-2q^2}{q^4}\frac{dK(q)^{n+1}}{dq^2} = \frac{1}{(4\pi)^2}\frac{2}{n+1}\underbrace{\left[-K(q)^{n+1}\right]_0^\infty}_{=1} \\ &= \boxed{\frac{1}{(4\pi)^2}\frac{2}{n+1}} \end{aligned}$$

Next we consider

$$J(n) \equiv \int_q \frac{\Delta(q)\,(1-K(q))^n}{q^4}$$

in $D = 4$, where n is a non-negative integer. Using the binomial expansion, we obtain

$$J(n) = \sum_{i=0}^{n}(-)^i\binom{n}{i}I(i) = \frac{2}{(4\pi)^2}\sum_{i=0}^{n}(-)^i\frac{1}{i+1}\binom{n}{i}$$

$$= \frac{2}{(4\pi)^2}\frac{1}{n+1}\underbrace{\sum_{i=0}^{n}(-)^i\binom{n+1}{i+1}}_{=1}$$

$$= \boxed{\frac{1}{(4\pi)^2}\frac{2}{n+1}}$$

Using this result, we can write down the one-loop vertices in the mass independent scheme as follows:



$$= -\frac{\lambda}{2}\left(-\frac{\Lambda^2}{2}\int_q \frac{\Delta(q)}{q^2} + m^2 \ln\frac{\Lambda}{\mu}\frac{2}{(4\pi)^2} + m^4\int_q \frac{1-K(q/\Lambda)}{q^4(q^2+m^2)}\right)$$



$$= \frac{\lambda^2}{(4\pi)^2}\ln\frac{\Lambda}{\mu}$$

$$+\frac{\lambda^2}{2}\int_q\left[\frac{1-K((q+p_1+p_2)/\Lambda)}{(q+p_1+p_2)^2+m^2}\frac{1-K(q/\Lambda)}{q^2+m^2} - \frac{(1-K(q/\Lambda))^2}{q^4}\right]$$

This implies

$$\left\{\begin{array}{rcl} b_2(\ln\Lambda/\mu,\lambda) &=& -\frac{\lambda}{(4\pi)^2}\ln\frac{\Lambda}{\mu}+\cdots \\ c_2(\ln\Lambda/\mu,\lambda) &=& \mathrm{O}\left(\lambda^2\right) \\ a_4(\ln\Lambda/\mu,\lambda) &=& \frac{3\lambda^2}{(4\pi)^2}\ln\frac{\Lambda}{\mu}+\cdots \end{array}\right.$$

2.7 The continuum limit in terms of a cutoff theory

Before diving into more detailed analysis of ERG, let us take a moment to reflect on what ERG gives us. In deriving ERG, we have made sure that the entire physics is kept untouched when we lower the cutoff Λ. No matter what Λ we use, we can always compute the same correlation functions. Let us give this statement a more concrete expression.

We recall that the two actions $S(\Lambda)$ and $S(\Lambda')$ on the same ERG trajectory are related as follows:

$$\left\{\begin{array}{rcl} \langle\phi(p)\phi(-p)\rangle_{S(\Lambda)} - \frac{K(p/\Lambda)}{p^2+m^2} &=& \frac{K(p/\Lambda)^2}{K(p/\Lambda')^2}\left(\langle\phi(p)\phi(-p)\rangle_{S(\Lambda')} - \frac{K(p/\Lambda')}{p^2+m^2}\right) \\ \langle\phi(p_1)\cdots\phi(p_{2n})\rangle_{S(\Lambda)} &=& \prod_{i=1}^{2n}\frac{K(p_i/\Lambda)}{K(p_i/\Lambda')}\cdot\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_{S(\Lambda')} \end{array}\right.$$

This can be rewritten as

$$\left\{\begin{array}{rcl} \frac{1}{K(p/\Lambda)^2}\langle\phi(p)\phi(-p)\rangle_{S(\Lambda)} + \frac{1-1/K(p/\Lambda)}{p^2+m^2} &=& \frac{1}{K(p/\Lambda')^2}\langle\phi(p)\phi(-p)\rangle_{S(\Lambda')} + \frac{1-1/K(p/\Lambda')}{p^2+m^2} \\ \prod_{i=1}^{2n}\frac{1}{K(p_i/\Lambda)}\cdot\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_{S(\Lambda)} &=& \prod_{i=1}^{2n}\frac{1}{K(p_i/\Lambda')}\cdot\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_{S(\Lambda')} \end{array}\right.$$

If the action corresponds to a continuum limit, we can take $\Lambda\to\infty$. Hence, the correlation functions in the continuum limit can be obtained from the action $S(\Lambda)$ with a finite cutoff:

$$\left\{\begin{array}{rcl} \langle\phi(p)\phi(-p)\rangle_\infty &=& \frac{1}{K(p/\Lambda)^2}\langle\phi(p)\phi(-p)\rangle_{S(\Lambda)} + \frac{1-1/K(p/\Lambda)}{p^2+m^2} \\ \langle\phi(p_1)\cdots\phi(p_{2n})\rangle_\infty &=& \prod_{i=1}^{2n}\frac{1}{K(p_i/\Lambda)}\cdot\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_{S(\Lambda)} \end{array}\right.$$

Here the left-hand sides are the correlation functions calculated with $S(\Lambda \to \infty)$. The right-hand sides are independent of Λ. It is surprising that the continuum physics can be obtained from an action with a finite cutoff, but we must accept it. There is a trade off, though. Our action $S(\Lambda)$ has an infinite number of interaction terms, and is highly complicated.

The above observation has a surprising consequence. If we expect the presence of a symmetry in the continuum limit, whether it is global or local, it should be present in the cutoff theory, too. Hence, for example, we expect to be able to construct gauge theories using a finite cutoff. We will discuss QED in some details, and sketch the construction of YM theories, both using a finite cutoff.

2.8 ERG for the 1PI $\Gamma[\Phi]$

This subsection is a digression from the main flow of the lectures. You can skip this part if you like. In the literature[4] the ERG differential equation for S_{int} is sometimes written for its 1PI part Γ_{int}. Roughly speaking, S_{int} consists of Feynman diagrams with elementary vertices and high-momentum propagators. These Feynman graphs are not necessarily 1PI, as we have seen in the case of $\mathcal{V}_4^{(1)}$ in the previous subsection. We wish to introduce 1PI vertices Γ_{int} so that S_{int} is obtained as the tree diagrams with high-momentum propagators and vertices Γ_{int}. The purpose of this subsection is to introduce Γ_{int} and ERG differential equation it satisfies. We will be sketchy since we will not discuss Γ_{int} in the remainder of the lectures.

Let us consider

$$\Gamma[\phi] \equiv -\frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2+m^2}{1-K(p/\Lambda)} + \Gamma_{\text{int}}[\phi]$$

We define its Legendre transform by

$$W[J] \equiv \Gamma[\phi] + \int_p \phi(-p)J(p)$$

where ϕ is determined in terms of J as

$$\frac{\delta\Gamma[\phi]}{\delta\phi(-p)} + J(p) = 0$$

This can be rewritten as

$$\phi(p) = \frac{1-K(p/\Lambda)}{p^2+m^2}\left(J(p) + \frac{\delta\Gamma_{\text{int}}[\phi]}{\delta\phi(-p)}\right)$$

In order for Γ_{int} to give the 1PI part of S_{int}, we must demand that

$$W[J] = \frac{1}{2}\int_p J(p)J(-p)\frac{1-K(p/\Lambda)}{p^2+m^2} + S_{\text{int}}\left[\frac{1-K(p/\Lambda)}{p^2+m^2}J(p)\right]$$



Figure 10: $\Gamma_{\rm int}$ is the 1PI part of $S_{\rm int}$.

(Fig. 10) By substituting

$$J(p) = \frac{p^2+m^2}{1-K(p/\Lambda)}\phi(p)$$

we obtain

$$W\left[\frac{p^2+m^2}{1-K(p/\Lambda)}\phi(p)\right] = \frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2+m^2}{1-K(p/\Lambda)} + S_{\rm int}[\phi]$$

On the other hand, from the definition of the Legendre transform, we obtain

$$W\left[\frac{p^2+m^2}{1-K(p/\Lambda)}\phi(p)\right] = \Gamma[\Phi] + \int_p \Phi(p)\frac{p^2+m^2}{1-K(p/\Lambda)}\phi(-p)$$

where

$$\phi(p) = \Phi(p) - \frac{1-K(p/\Lambda)}{p^2+m^2}\frac{\delta\Gamma_{\rm int}[\Phi]}{\delta\Phi(-p)}$$

Hence, we obtain the following relation between $\Gamma_{\rm int}[\Phi]$ and $S_{\rm int}[\phi]$:

$$\begin{aligned}&-\frac{1}{2}\int_p \Phi(p)\Phi(-p)\frac{p^2+m^2}{1-K(p/\Lambda)} + \Gamma_{\rm int}[\Phi]\\ &\quad= \frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2+m^2}{1-K(p/\Lambda)} + S_{\rm int}[\phi] - \int_p \phi(p)\frac{p^2+m^2}{1-K(p/\Lambda)}\Phi(-p)\end{aligned}$$

It is a simple algebra to rewrite this as

$$\Gamma_{\rm int}[\Phi] = \frac{1}{2}\int_p \left(\phi(p)-\Phi(p)\right)\left(\phi(-p)-\Phi(-p)\right)\frac{p^2+m^2}{1-K(p/\Lambda)} + S_{\rm int}[\phi]$$

Now, we can reverse the direction, and obtain Γ by Legendre transforming W:

$$\Gamma[\Phi] = W[J] - \int_p \Phi(-p)J(p)$$

---

[4] For example, see [13].

where

$$\Phi(p) = \frac{\delta W[J]}{\delta J(-p)}$$

By the substitution $J(p) = \frac{p^2+m^2}{1-K(p/\Lambda)}\,\phi(p)$, this gives

$$\Phi(p) = \phi(p) + \frac{1-K(p/\Lambda)}{p^2+m^2}\frac{\delta S_{\mathrm{int}}[\phi]}{\delta\phi(-p)}$$

Comparing this with the relation between ϕ and Φ obtained above, we obtain

$$\frac{\delta\Gamma_{\mathrm{int}}[\Phi]}{\delta\Phi(-p)} = \frac{\delta S_{\mathrm{int}}[\phi]}{\delta\phi(-p)}$$

Now that we have an explicit relation between Γ_{int} and S_{int}, we can derive the ERG differential equation of Γ_{int} from that of S_{int}. The derivation is straightforward, and we only state the result:

$$-\Lambda\frac{\partial}{\partial\Lambda}\Gamma_{\mathrm{int}}[\Phi] = \frac{1}{2}\int_q \frac{\Delta(q/\Lambda)}{q^2+m^2}\frac{\delta^2 S_{\mathrm{int}}[\phi]}{\delta\phi(-q)\delta\phi(q)}$$

By differentiating the relation obtained above between the first order functional derivatives of Γ_{int} and S_{int}, we obtain

$$\frac{\delta^2 S_{\mathrm{int}}}{\delta\phi(q)\delta\phi(-q)} = \int_p \frac{\delta\Phi(p)}{\delta\phi(q)}\frac{\delta^2\Gamma_{\mathrm{int}}[\Phi]}{\delta\Phi(p)\delta\Phi(-q)}$$

where $\frac{\delta\Phi(p)}{\delta\phi(q)}$ is the inverse of

$$\frac{\delta\phi(q)}{\delta\Phi(p)} = (2\pi)^4\delta^{(4)}(p-q) - \frac{1-K(q/\Lambda)}{q^2+m^2}\frac{\delta^2\Gamma_{\mathrm{int}}[\Phi]}{\delta\Phi(p)\delta\Phi(-q)}$$

Hence, it is given by the geometric series

$$\begin{aligned}\frac{\delta\Phi(p)}{\delta\phi(q)} &= (2\pi)^4\delta^{(4)}(p-q) + \frac{1-K(p/\Lambda)}{p^2+m^2}\frac{\delta^2\Gamma_{\mathrm{int}}[\Phi]}{\delta\Phi(-p)\delta\Phi(q)} \\ &\qquad + \int_r \frac{1-K(p/\Lambda)}{p^2+m^2}\frac{\delta^2\Gamma_{\mathrm{int}}[\Phi]}{\delta\Phi(-p)\delta\Phi(r)}\frac{1-K(r/\Lambda)}{r^2+m^2}\frac{\delta^2\Gamma_{\mathrm{int}}}{\delta\Phi(-r)\delta\Phi(q)} + \cdots\end{aligned}$$

Therefore, we obtain

$$\frac{\delta^2 S_{\mathrm{int}}}{\delta\phi(q)\delta\phi(-q)} = \frac{\delta^2\Gamma_{\mathrm{int}}}{\delta\Phi(q)\delta\Phi(-q)} + \int_r \frac{\delta^2\Gamma_{\mathrm{int}}}{\delta\Phi(q)\delta\Phi(-r)}\frac{1-K(r/\Lambda)}{r^2+m^2}\frac{\delta^2\Gamma_{\mathrm{int}}}{\delta\Phi(r)\delta\Phi(-q)} + \cdots$$

Thus, we can express this ERG equation graphically as



This final result is actually easier to obtain diagrammatically.
HW#4: From the above diagrammatic ERG equation for Γ_{int}, derive the ERG equation for S_{int} diagrammatically.

2.9 Composite operators

The purpose of this subsection is to introduce composite operators which play crucial roles in understanding some important properties of ERG such as its relation to beta functions and anomalous dimensions. We take $D = 4$ for concreteness.

We call

$$\begin{aligned}\mathcal{O}(p) \quad &\equiv \quad \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1,\cdots,p_{2n}} \phi(p_1)\cdots\phi(p_{2n})\mathcal{O}_{2n}(\Lambda; p_1, \cdots, p_{2n}) \\ &\qquad \times (2\pi)^4 \delta^{(4)}(p_1 + \cdots + p_{2n} - p)\end{aligned}$$

a **composite operator** [5] if it satisfies the ERG differential equation

$$-\Lambda \frac{\partial}{\partial \Lambda} \mathcal{O}(p) = \mathcal{D} \cdot \mathcal{O}(p)$$

where $\mathcal{D}$ is the differential operator

$$\mathcal{D} \equiv \int_q \frac{\Delta(q/\Lambda)}{q^2 + m^2} \left(\frac{\delta S_{\mathrm{int}}}{\delta\phi(-q)} \frac{\delta}{\delta\phi(-q)} + \frac{1}{2} \frac{\delta^2}{\delta\phi(q)\delta\phi(-q)} \right)$$

We call $\mathcal{O}$ an operator of dimension d, if it satisfies the asymptotic conditions

$$\mathcal{O}_{2n \geq d+2}(\Lambda; p_1, \cdots, p_{2n}) \overset{\Lambda\to\infty}{\longrightarrow} 0$$

Hence, an operator of dimension 2 has $\mathcal{O}_{2n\geq 4} \to 0$, and that of dimension 4 has $\mathcal{O}_{2n\geq 6} \to 0$.

It is easy to understand where the above definition of composite operators comes from. Let us consider a modified action

$$S'(\Lambda) \equiv S(\Lambda) + \int_p \epsilon(-p)\mathcal{O}(p)$$

where ϵ is an infinitesimal source. The definition guarantees that S' satisfies the Polchinski differential equation if S satisfies it.

HW#5: Show that the ERG differential equation for the composite operator $\mathcal{O}$ can be rewritten as

$$-\Lambda \frac{\partial}{\partial \Lambda} \mathcal{O} = \int_q \frac{\Delta(q/\Lambda)}{q^2 + m^2} \left(\frac{q^2 + m^2}{K(q/\Lambda)} \phi(q) \frac{\delta}{\delta\phi(q)} + \frac{\delta S}{\delta\phi(-q)} \frac{\delta}{\delta\phi(q)} + \frac{1}{2} \frac{\delta^2}{\delta\phi(q)\delta\phi(-q)} \right) \mathcal{O}$$

Let us recall the following Λ dependence of the correlation functions:

$$\begin{aligned}\langle\phi(p)\phi(-p)\rangle_{S(\Lambda)} - \tfrac{K(p/\Lambda)}{p^2+m^2} \quad &= \quad \tfrac{K(p/\Lambda)^2}{K(p/\Lambda')^2} \left(\langle\phi(p)\phi(-p)\rangle_{S(\Lambda')} - \tfrac{K(p/\Lambda')}{p^2+m^2} \right) \\ \langle\phi(p_1)\cdots\phi(p_{2n})\rangle_{S(\Lambda)} \quad &= \quad \textstyle\prod_{i=1}^{2n} \tfrac{K(p_i/\Lambda)}{K(p_i/\Lambda')} \cdot \langle\phi(p_1)\cdots\phi(p_{2n})\rangle_{S(\Lambda')} \quad (n > 1)\end{aligned}$$

---

[5] Here we assume $\mathcal{O}$ is even in ϕ. We can also define $\mathcal{O}$ odd in ϕ. For example, $\frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}$ considered below is odd in ϕ.

Substituting S' into S and taking the part proportional to $\epsilon(-p)$, we obtain

$$\langle \phi(p_1)\cdots\phi(p_{2n})\mathcal{O}(p)\rangle_{S(\Lambda)} = \prod_{i=1}^{2n} \frac{K(p_i/\Lambda)}{K(p_i/\Lambda')} \cdot \langle \phi(p_1)\cdots\phi(p_{2n})\mathcal{O}(p)\rangle_{S(\Lambda')}$$

for any $n \geq 1$. Equivalently, this gives

$$\langle \phi(p_1)\cdots\phi(p_{2n})\mathcal{O}(p)\rangle_{\infty} = \prod_{i=1}^{2n} \frac{1}{K(p_i/\Lambda)} \cdot \langle \phi(p_1)\cdots\phi(p_{2n})\mathcal{O}(p)\rangle_{S(\Lambda)}$$

We can actually reverse the direction, and use either of the above two relations as the defining property of a composite operator.

The simplest example of a composite operator is $\mathcal{O}(p) \equiv \frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}$. To show this, we simply differentiate the ERG differential equation

$$-\Lambda\frac{\partial}{\partial\Lambda}S_{\mathrm{int}} = \int_q \frac{\Delta(q/\Lambda)}{q^2+m^2}\frac{1}{2}\left\{\frac{\delta S_{\mathrm{int}}}{\delta\phi(q)}\frac{\delta S_{\mathrm{int}}}{\delta\phi(-q)} + \frac{\delta^2 S_{\mathrm{int}}}{\delta\phi(q)\delta\phi(-q)}\right\}$$

with respect to $\phi(-p)$. The asymptotic behavior of S_{int} (in the mass independent scheme) translates into the following asymptotic behavior of $\frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}$:

$$\begin{array}{rcl} \mathcal{O}_1(p) & \longrightarrow & a_2(\ln\Lambda/\mu,\lambda) + m^2 b_2(\ln\Lambda/\mu,\lambda) + p^2 c_2(\ln\Lambda/\mu,\lambda) \\ \mathcal{O}_3(p_1,p_2,p_3) & \longrightarrow & a_4(\ln\Lambda/\mu,\lambda) \\ \mathcal{O}_{n\geq 5}(p_1,\cdots,p_n) & \longrightarrow & 0 \end{array}$$

Hence, $\frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}$ is dimension 3. Since $\frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}$ is a composite operator, the correlation functions

$$\left\langle \frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}\phi(p_1)\cdots\phi(p_{2n-1})\right\rangle_{\infty} = \left\langle \frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}\phi(p_1)\cdots\phi(p_{2n-1})\right\rangle_{S(\Lambda)} \cdot \prod_{i=1}^{2n-1}\frac{1}{K(p_i/\Lambda)}$$

are independent of Λ.

Another example is

$$\Phi(p) \equiv \phi(p) + \frac{1-K(p/\Lambda)}{p^2+m^2}\frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}$$

which corresponds to an elementary field $\phi(p)$. To show this, we can again take the Λ derivative of Φ:

$$-\Lambda\frac{\partial\Phi(p)}{\partial\Lambda} = \frac{\Delta(p/\Lambda)}{p^2+m^2}\frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)} + \mathcal{D}\cdot\frac{1-K(p/\Lambda)}{p^2+m^2}\frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}$$

Since

$$\mathcal{D}\cdot\phi(p) = \frac{\Delta(p/\Lambda)}{p^2+m^2}\frac{\delta S_{\mathrm{int}}}{\delta\phi(-p)}$$

we obtain

$$-\Lambda\frac{\partial\Phi(p)}{\partial\Lambda} = \mathcal{D}\cdot\Phi(p)$$

HW#5: Show that

$$K(p/\Lambda)\frac{\delta S}{\delta\phi(-p)} = -(p^2+m^2)\Phi(p) + \frac{\delta S_{\rm int}}{\delta\phi(-p)}$$

and hence it is a composite operator.

This is an operator corresponding to the equation of motion. Since

$$\begin{aligned}\frac{\delta}{\delta\phi(-p)}\left(\phi(p_1)\cdots\phi(p_n)\mathrm{e}^S\right) &= \frac{\delta S}{\delta\phi(-p)}\phi(p_1)\cdots\phi(p_n)\mathrm{e}^S\\ &+\sum_{i=1}^n (2\pi)^4\delta^{(4)}(p-p_i)\,\phi(p_1)\cdots\widehat{\phi(p_i)}\cdots\phi(p_n)\mathrm{e}^S\end{aligned}$$

is a total derivative, its functional integral vanishes. Hence, integrating over ϕ, we obtain

$$\left\langle \frac{\delta S}{\delta\phi(-p)}\phi(-p)\right\rangle_{S(\Lambda)} = -1$$

and for $n>1$

$$\left\langle \frac{\delta S}{\delta\phi(-p)}\phi(p_1)\cdots\phi(p_n)\right\rangle_{S(\Lambda)} = 0$$

if we take the connected part. Thus, in the continuum limit the equations of motion are given by

$$\left\{\begin{array}{rcll} \left\langle K(p/\Lambda)\frac{\delta S}{\delta\phi(-p)}\phi(-p)\right\rangle_\infty &=& -1 & \\ \left\langle K(p/\Lambda)\frac{\delta S}{\delta\phi(-p)}\phi(p_1)\cdots\phi(p_{2n-1})\right\rangle_\infty &=& 0 & (n>1)\end{array}\right.$$

2.10 Composite operators given in terms of the action

As for the action itself, the ERG differential equation of a composite operator does not determine the operator uniquely. We must supply appropriate asymptotic conditions. For later conveniences, we consider dimension 2 and 4 scalar composite operators. We adopt the mass independent scheme for simplicity.

Dimension 2 operator

There is a unique dimension 2 operator $\mathcal{O}$ of zero momentum, specified by the following asymptotic conditions[6]:

$$\left\{\begin{array}{ccl} \mathcal{O}_2(\Lambda;p,-p) & \longrightarrow & a_2(\ln\Lambda/\mu;\lambda)\\ \mathcal{O}_{2n\ge4}(\Lambda;p_1,\cdots,p_{2n}) & \longrightarrow & 0\end{array}\right.$$

where

$$a_2(0;\lambda)=1$$

We call this operator $\left[\frac{1}{2}\phi^2\right]$.

---

[6] This a_2 is not the same as the a_2 that gives the asymptotic behavior of the vertex $\mathcal{V}_2$.

Dimension 4 operators

Dimension 4 operators $\mathcal{O}$ of zero momentum are specified by the following asymptotic conditions:

$$\left\{\begin{array}{ccl} \mathcal{O}_2(\Lambda; p, -p) & \longrightarrow & \Lambda^2 a_2(\ln \Lambda/\mu; \lambda) + m^2 b_2(\ln \Lambda/\mu; \lambda) + p^2 c_2(\ln \Lambda/\mu; \lambda) \\ \mathcal{O}_4(\Lambda; p_1, \cdots, p_4) & \longrightarrow & a_4(\ln \Lambda/\mu; \lambda) \\ \mathcal{O}_{2n\geq 6}(\Lambda; p_1, \cdots, p_{2n}) & \longrightarrow & 0 \end{array}\right.$$

There are two linearly independent operators:

1. $\left[\frac{1}{4!}\phi^4\right]$, satisfying
$$a_4(0;\lambda) = 1, \quad b_2(0;\lambda) = c_2(0;\lambda) = 0$$
2. $\left[\frac{1}{2}(\partial_\mu \phi)^2\right]$, satisfying
$$c_2(0;\lambda) = 1, \quad b_2(0;\lambda) = a_4(0;\lambda) = 0$$

We can regard $m^2\left[\frac{1}{2}\phi^2\right]$ as the third dimension 4 operator satisfying

$$b_2(0;\lambda) = 1, \quad c_2(0;\lambda) = a_4(0;\lambda) = 0$$

Composite operators given in terms of S

The purpose of the remaining part of this subsection is to show that the three dimension 4 scalar operators

$$m^2\left[\frac{1}{2}\phi^2\right], \quad \left[\frac{1}{2}\left(\partial_\mu \phi\right)^2\right], \quad \left[\frac{1}{4!}\phi^4\right]$$

are obtained in terms of the action S.

As a preparation, we consider the following:

$$\left\{\begin{array}{ccl} \mathcal{O}_1[f] & \equiv & \int_p f(p)\phi(p)\frac{\delta S}{\delta\phi(p)} \\ \mathcal{O}_2[C] & \equiv & \int_p C(p^2)\frac{1}{2}\left\{\frac{\delta S}{\delta\phi(p)}\frac{\delta S}{\delta\phi(-p)} + \frac{\delta^2 S}{\delta\phi(p)\delta\phi(-p)}\right\} \end{array}\right.$$

Neither of them is a composite operator by itself, but their correlation functions can be easily computed.

1. $\mathcal{O}_1[f]$ — Except for an inessential additive constant[7], $\mathcal{O}_1[f]$ is a total derivative
$$\mathcal{O}_1[f]\mathrm{e}^S = \int_p f(p)\frac{\delta}{\delta\phi(p)}\left(\phi(p)\mathrm{e}^S\right)$$
Hence, multiplying this by $\phi(p_1)\cdots\phi(p_{2n})$ and integrating over ϕ by parts, we obtain
$$\langle \mathcal{O}_1[f]\phi(p_1)\cdots\phi(p_{2n})\rangle_S = -\left(\sum_{i=1}^{2n} f(p_i)\right)\cdot\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_S$$

---

[7] The additive constant does not contribute to the connected part.

2. $\mathcal{O}_2[C]$ — This is also a total derivative:

$$\mathcal{O}_2[C]\mathrm{e}^S = \int_p C(p^2)\frac{1}{2}\frac{\delta^2}{\delta\phi(p)\delta\phi(-p)}\,\mathrm{e}^S$$

   Again, integrating over ϕ by parts, we obtain

$$\left\{\begin{array}{rcl} \langle\mathcal{O}_2[C]\phi(p)\phi(-p)\rangle_S &=& C(p^2) \\ \langle\mathcal{O}_2[C]\phi(p_1)\cdots\phi(p_{2n})\rangle_S &=& 0 \quad (n>1) \end{array}\right.$$

It is interesting to note that the right-hand side of the Polchinski differential ERG equation is written as

$$\int_q \frac{\Delta(q/\Lambda)}{q^2+m^2}\left[\frac{q^2+m^2}{K(q/\Lambda)}\phi(q)\frac{\delta S}{\delta\phi(q)} + \frac{1}{2}\left\{\frac{\delta S}{\delta\phi(q)}\frac{\delta S}{\delta\phi(-q)} + \frac{\delta^2 S}{\delta\phi(q)\delta\phi(-q)}\right\}\right]$$
$$= \mathcal{O}_1[f] + \mathcal{O}_2[C]$$

where

$$\left\{\begin{array}{rcl} f(q) &=& \frac{\Delta(q/\Lambda)}{K(q/\Lambda)} \\ C(q^2) &=& \frac{\Delta(q/\Lambda)}{q^2+m^2} \end{array}\right.$$

Hence, we reproduce

$$\left\{\begin{array}{rcl} -\Lambda\frac{\partial}{\partial\Lambda}\langle\phi(p)\phi(-p)\rangle_S &=& \frac{\Delta(p/\Lambda)}{p^2+m^2} - 2\frac{\Delta(p/\Lambda)}{K(p/\Lambda)}\langle\phi(p)\phi(-p)\rangle_S \\ -\Lambda\frac{\partial}{\partial\Lambda}\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_S &=& -\sum_{i=1}^{2n}\frac{\Delta(p_i/\Lambda)}{K(p_i/\Lambda)}\cdot\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_S \end{array}\right.$$

With the above preparation, we can construct three special operators $\mathcal{O}_m, \mathcal{O}_\lambda$, and $\mathcal{N}$.

1. $\mathcal{O}_m$ that generates the change of correlation functions with respect to m^2 — by differentiating the correlation functions with respect to m^2, we obtain

$$-\partial_{m^2}\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_S = \langle(-\partial_{m^2}S)\,\phi(p_1)\cdots\phi(p_{2n})\rangle_S \quad (n\ge 1)$$

   Hence, for $n>1$, we obtain

$$-\partial_{m^2}\langle\phi(p_1)\cdots\phi(p_{2n})\rangle_\infty = \langle(-\partial_{m^2}S)\,\phi(p_1)\cdots\phi(p_{2n})\rangle_S\prod_{i=1}^{2n}\frac{1}{K(p_i/\Lambda)}$$

   The case $n=1$ needs extra care. Since

$$\langle\phi(p)\phi(-p)\rangle_\infty = \frac{1-1/K(p/\Lambda)}{p^2+m^2} + \frac{1}{K(p/\Lambda)^2}\langle\phi(p)\phi(-p)\rangle_S$$

   we obtain

$$-\partial_{m^2}\langle\phi(p)\phi(-p)\rangle_\infty = \frac{1-1/K(p/\Lambda)}{(p^2+m^2)^2} + \frac{1}{K(p/\Lambda)^2}\langle(-\partial_{m^2}S)\,\phi(p)\phi(-p)\rangle_S$$
$$= \left\{-\frac{K(p/\Lambda)\,(1-K(p/\Lambda))}{(p^2+m^2)^2} + \langle(-\partial_{m^2}S)\,\phi(p)\phi(-p)\rangle_S\right\}\frac{1}{K(p/\Lambda)^2}$$

Thus, we obtain

$$-\partial_{m^2} \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty = \langle \mathcal{O}_m \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty \quad (n \geq 1)$$

for the composite operator

$$\boxed{\mathcal{O}_m \equiv \mathcal{O}_2[C] - \partial_{m^2} S}$$

where

$$C(p^2) \equiv - \frac{K(p/\Lambda)\,(1-K(p/\Lambda))}{(p^2+m^2)^2}$$

2. $\mathcal{O}_\lambda$ that generates the change with respect to λ — by differentiating the correlation functions with respect to λ, we obtain

$$-\partial_\lambda \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_S = \langle (-\partial_\lambda S)\, \phi(p_1) \cdots \phi(p_{2n}) \rangle_S \quad (n \geq 1)$$

   Hence,

$$\boxed{\mathcal{O}_\lambda \equiv -\partial_\lambda S}$$

   is a composite operator.

3. $\mathcal{N}$ that generates the change of normalization of ϕ — we wish to construct a composite operator $\mathcal{N}$ that counts the number of fields:

$$\langle \mathcal{N} \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty = 2n \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty \quad (n \geq 1)$$

   The starting point is $\mathcal{O}_1[f]$, where $f = -1$, which satisfies

$$\langle \mathcal{O}_1[-1] \phi(p_1) \cdots \phi(p_{2n}) \rangle_S = 2n \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_S \quad (n \geq 1)$$

   Again, the case $n = 1$ needs extra care. Since

$$2 \langle \phi(p)\phi(-p) \rangle_\infty = 2 \left(- \frac{K(p/\Lambda)\,(1-K(p/\Lambda))}{p^2+m^2} + \langle \phi(p)\phi(-p) \rangle_S \right) \frac{1}{K(p/\Lambda)^2}$$

   we obtain

$$\boxed{\mathcal{N} = \mathcal{O}_2[C] + \mathcal{O}_1[-1]}$$

   where

$$C(p^2) = -2 \frac{K(p/\Lambda)\,(1-K(p/\Lambda))}{p^2+m^2}$$

To summarize, we have constructed three dimension 4 composite operators $\mathcal{O}_m, \mathcal{O}_\lambda, \mathcal{N}$ with the following properties:

$$\boxed{\begin{array}{rcl} \langle \mathcal{O}_m \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty &=& -\partial_{m^2} \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty \\ \langle \mathcal{O}_\lambda \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty &=& -\partial_\lambda \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty \\ \langle \mathcal{N} \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty &=& 2n \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty \end{array}}$$

where these operators are given in terms of the action as

$$\left\{\begin{array}{rcl} \mathcal{O}_m & \equiv & -\partial_{m^2} S - \int_q \frac{K(q/\Lambda)(1-K(q/\Lambda))}{(q^2+m^2)^2} \frac{1}{2} \left\{ \frac{\delta S}{\delta\phi(q)} \frac{\delta S}{\delta\phi(-q)} + \frac{\delta^2 S}{\delta\phi(q)\delta\phi(-q)} \right\} \\ \mathcal{O}_\lambda & \equiv & -\partial_\lambda S \\ \mathcal{N} & \equiv & -\int_q \phi(q) \frac{\delta S}{\delta\phi(q)} - \int_q \frac{K(q/\Lambda)(1-K(q/\Lambda))}{q^2+m^2} \left\{ \frac{\delta S}{\delta\phi(q)} \frac{\delta S}{\delta\phi(-q)} + \frac{\delta^2 S}{\delta\phi(q)\delta\phi(-q)} \right\} \end{array}\right.$$

These operators are obtained as linear combinations of $\left[\frac{\phi^2}{2}\right]$, $\left[\frac{\phi^4}{4!}\right]$, and $\left[\frac{1}{2}(\partial_\mu\phi)^2\right]$. To see this, we must examine the asymptotic behaviors as $\Lambda \to \infty$. We first introduce the following notation[8]:

$$\begin{array}{rcl} \mathcal{V}_4(\Lambda; q, -q, 0, 0) & \longrightarrow & a_4(\ln\Lambda/\mu; q/\Lambda) + \frac{m^2}{\Lambda^2} b_4(\ln\Lambda/\mu; q/\Lambda) + \cdots \\ \mathcal{V}_6(\Lambda; q, -q, 0, \cdots, 0) & \longrightarrow & \frac{1}{\Lambda^2} a_6(\ln\Lambda/\mu; q/\Lambda) + \cdots \end{array}$$

Then, using the mass independent scheme, it is straightforward to derive the following results[9]:

$$\boxed{\begin{array}{rcl} \mathcal{O}_m & = & x_m \left[\frac{1}{2}\phi^2\right] \\ \mathcal{O}_\lambda & = & \left[\frac{1}{4!}\phi^4\right] \\ \mathcal{N} & = & x_{\mathcal{N}} m^2 \left[\frac{1}{2}\phi^2\right] + y_{\mathcal{N}} \left[\frac{1}{4!}\phi^4\right] + z_{\mathcal{N}} \left[\frac{1}{2}(\partial_\mu\phi)^2\right] \end{array}}$$

where

$$\left\{\begin{array}{rcl} x_m & = & 1 - \frac{1}{2}\int_q \frac{K(q)(1-K(q))}{q^4} a_4(0;q) \\ x_{\mathcal{N}} & = & 2 + \int_q K(q)(1-K(q)) \left(-\frac{b_4(0;q)}{q^2} + \frac{a_4(0;q)}{q^4} \right) \\ y_{\mathcal{N}} & = & -4\lambda - \int_q \frac{K(q)(1-K(q))}{q^2} a_6(0;q) \\ z_{\mathcal{N}} & = & 2 - \int_q \frac{K(q)(1-K(q))}{q^2} c_4(0;q) \end{array}\right.$$

where c_4 is defined by

$$\frac{1}{\Lambda^2} c_4(\ln\lambda/\mu; q/\Lambda) \equiv \frac{\partial}{\partial p^2} \mathcal{V}_4(\Lambda; p, -p, q, -q)\Big|_{m^2=p^2=0}$$

where the angular average over q_μ is taken.

To conclude this subsection, we invert the relation of $\mathcal{O}_m, \mathcal{O}_\lambda, \mathcal{N}$ to the mass independent composite operators to obtain

$$\boxed{\begin{array}{rcl} \left[\frac{1}{2}\phi^2\right] & = & \frac{1}{x_m}\mathcal{O}_m \\ \left[\frac{1}{4!}\phi^4\right] & = & \mathcal{O}_\lambda \\ \left[\frac{1}{2}(\partial_\mu\phi)^2\right] & = & \frac{1}{z_{\mathcal{N}}} \left(\mathcal{N} - \frac{x_{\mathcal{N}}}{x_m}\mathcal{O}_m - y_{\mathcal{N}}\mathcal{O}_\lambda \right) \end{array}}$$

---

[8] These are expansions in powers of $\frac{1}{\Lambda^2}$

[9] It is “straightforward” but some work is required.

2.11 Beta functions and anomalous dimensions

To be concrete, we adopt the mass independent scheme to define the action $S(\Lambda; m^2, \lambda; \mu)$. The action depends on three parameters: m^2, λ, and μ. Here, μ is an arbitrary momentum scale that we have used to specify a unique solution to the ERG differential equation. We can physically interpret m^2 and λ as the mass and interaction parameters of the theory defined at the scale μ.

Let us suppose we change the scale μ. In order to keep the physics (or equivalently correlation functions) intact, we must change m^2 and λ accordingly. This change is the conventional RG transformation of the running parameters m^2 and λ.

We first observe that

$$\Psi \equiv -\mu \frac{\partial}{\partial \mu} S(\Lambda; m^2, \lambda; \mu)$$

is a composite operator; for $\mu \neq \mu'$ both $S(\Lambda; m^2, \lambda; \mu)$ and $S(\Lambda; m^2, \lambda; \mu')$ satisfy the same ERG differential equation. From its asymptotic behaviors, we find Ψ a dimension 4 operator, and hence it is given as a linear combination of $m^2 \mathcal{O}_m$, $\mathcal{O}_\lambda$, and $\mathcal{N}$. To determine the coefficients, we must examine the asymptotic behaviors of Ψ in details.

We first recall the asymptotic behavior of $S_{\rm int}(\Lambda)$:

$$\left\{ \begin{array}{rcl} \mathcal{V}_2(\Lambda; p, -p) & \longrightarrow & \Lambda^2 a_2(\ln \Lambda/\mu, \lambda) + m^2 b_2(\ln \Lambda/\mu, \lambda) + p^2 c_2(\ln \Lambda/\mu, \lambda) \\ \mathcal{V}_4(\Lambda; p_1, \cdots, p_4) & \longrightarrow & a_4(\ln \Lambda/\mu, \lambda) \end{array} \right.$$

Hence, the asymptotic behavior of $\Psi(\Lambda)$ is given by

$$\left\{ \begin{array}{rcl} \Psi_2(\Lambda; p, -p) & \longrightarrow & \Lambda^2 \dot{a}_2(\ln \Lambda/\mu, \lambda) + m^2 \dot{b}_2(\ln \Lambda/\mu, \lambda) + p^2 \dot{c}_2(\ln \Lambda/\mu, \lambda) \\ \Psi_4(\Lambda; p_1, \cdots, p_4) & \longrightarrow & \dot{a}_4(\ln \Lambda/\mu, \lambda) \end{array} \right.$$

where we have defined

$$\left\{ \begin{array}{rcccl} \dot{a}_2(\ln \Lambda/\mu, \lambda) & \equiv & -\mu \frac{\partial}{\partial \mu} a_2(\ln \Lambda/\mu, \lambda) & = & \Lambda \frac{\partial}{\partial \Lambda} a_2(\ln \Lambda/\mu, \lambda) \\ \dot{b}_2(\ln \Lambda/\mu, \lambda) & \equiv & -\mu \frac{\partial}{\partial \mu} b_2(\ln \Lambda/\mu, \lambda) & = & \Lambda \frac{\partial}{\partial \Lambda} b_2(\ln \Lambda/\mu, \lambda) \\ \dot{c}_2(\ln \Lambda/\mu, \lambda) & \equiv & -\mu \frac{\partial}{\partial \mu} c_2(\ln \Lambda/\mu, \lambda) & = & \Lambda \frac{\partial}{\partial \Lambda} c_2(\ln \Lambda/\mu, \lambda) \\ \dot{a}_4(\ln \Lambda/\mu, \lambda) & \equiv & -\mu \frac{\partial}{\partial \mu} a_4(\ln \Lambda/\mu, \lambda) & = & \Lambda \frac{\partial}{\partial \Lambda} a_4(\ln \Lambda/\mu, \lambda) \end{array} \right.$$

We denote the values of these at $\Lambda = \mu$ using the same notation as follows:

$$\dot{a}_2(\lambda) \equiv \dot{a}_2(0, \lambda),\ \dot{b}_2(\lambda) \equiv \dot{b}_2(0, \lambda),\ \cdots$$

Then, comparing the asymptotic behaviors of Ψ with those of $\left[\frac{1}{2}\phi^2\right]$, $\left[\frac{1}{4!}\phi^4\right]$, and $\left[\frac{1}{2}\left(\partial_\mu \phi\right)^2\right]$, we obtain

$$\Psi = \dot{b}_2 m^2 \left[\frac{1}{2}\phi^2\right] + \dot{c}_2 \left[\frac{1}{2}\left(\partial_\mu \phi\right)^2\right] + \dot{a}_4 \left[\frac{1}{4!}\phi^4\right]$$

We can rewrite this using $\mathcal{O}_m, \mathcal{O}_\lambda$, and $\mathcal{N}$ as

$$\begin{aligned} \Psi &= \frac{\dot{b}_2}{x_m} m^2 \mathcal{O}_m + \dot{a}_4 \mathcal{O}_\lambda + \frac{\dot{c}_2}{z_{\mathcal{N}}} \left(\mathcal{N} - \frac{x_{\mathcal{N}}}{x_m} m^2 \mathcal{O}_m - y_{\mathcal{N}} \mathcal{O}_\lambda \right) \\ &= \frac{1}{x_m} \left(\dot{b}_2 - \frac{x_{\mathcal{N}}}{z_{\mathcal{N}}} \dot{c}_2 \right) m^2 \mathcal{O}_m + \left(\dot{a}_4 - \frac{y_{\mathcal{N}}}{z_{\mathcal{N}}} \dot{c}_2 \right) \mathcal{O}_\lambda + \frac{\dot{c}_2}{z_{\mathcal{N}}} \mathcal{N} \end{aligned}$$

Finally, by defining the following functions of λ

$$\left\{ \begin{array}{rcl} \beta_m(\lambda) &\equiv& \frac{1}{x_m} \left(\dot{b}_2 - \frac{x_{\mathcal{N}}}{z_{\mathcal{N}}} \dot{c}_2 \right) \\ \beta(\lambda) &\equiv& \dot{a}_4 - \frac{y_{\mathcal{N}}}{z_{\mathcal{N}}} \dot{c}_2 \\ \gamma(\lambda) &\equiv& \frac{\dot{c}_2}{z_{\mathcal{N}}} \end{array} \right.$$

we obtain

$$\boxed{\Psi = \beta_m m^2 \mathcal{O}_m + \beta \mathcal{O}_\lambda + \gamma \mathcal{N}}$$

This implies

$$\begin{aligned} -\mu \frac{\partial}{\partial \mu} \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty &= \langle \Psi \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty \\ &= \left(-\beta_m m^2 \partial_{m^2} - \beta \partial_\lambda + 2n\gamma \right) \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty \end{aligned}$$

Hence, we obtain the standard RG equation[10]

$$\boxed{\left(-\mu \frac{\partial}{\partial \mu} + \beta_m m^2 \partial_{m^2} + \beta \partial_\lambda - 2n\gamma \right) \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_\infty = 0}$$

This result was obtained in [15].

One-loop calculations

At one-loop, we have

$$\dot{b}_2 = -\frac{\lambda}{(4\pi)^2}, \quad \dot{c}_2 = 0, \quad \dot{a}_4 = \frac{3\lambda^2}{(4\pi)^2}$$

and to the lowest order

$$x_m = 1, \quad x_{\mathcal{N}} = 2, \quad y_{\mathcal{N}} = -4\lambda, \quad z_{\mathcal{N}} = 2$$

Hence, we obtain the familiar results:

$$\beta_m = -\frac{\lambda}{(4\pi)^2}, \quad \beta = \frac{3\lambda^2}{(4\pi)^2}, \quad \gamma = 0$$

HW#6: Check this result.

---

[10] Our β, β_m, γ have the opposite sign convention than the one usually used.

2.12 Universality

Universality is an important concept in renormalization theory. To construct a continuum limit, there are always more than one way. The independence of the continuum limit on the particular method of construction is called universality. For example, if we use a lattice to construct a continuum theory, the limit should not depend on what kind of lattice, whether square or cubic, we use.

In the following we examine universality in two restricted contexts. First, we wish to show that the continuum limit (i.e., the correlation functions with the suffix ∞) does not depend on the asymptotic conditions we use to select a particular solution of the ERG differential equation. Second, we wish to show that the continuum limit does not depend on the choice of a cutoff function K.

Scheme dependence

Given the ERG differential equation

$$-\Lambda \frac{\partial}{\partial \Lambda} S(\Lambda) = \int_q \frac{\Delta(q/\Lambda)}{q^2+m^2} \frac{1}{2} \left\{ \frac{\delta S}{\delta\phi(q)} \frac{\delta S}{\delta\phi(-q)} + \frac{\delta^2 S}{\delta\phi(q)\delta\phi(-q)} \right\}$$

the solution which corresponds to a renormalized theory has the following asymptotic behaviors for large Λ:

$$\left\{ \begin{array}{ccl} \mathcal{V}_2(\Lambda; p, -p) & \longrightarrow & \Lambda^2 a_2(m^2/\mu^2, \ln \Lambda/\mu, \lambda) \\ & & + m^2 b_2(m^2/\mu^2, \ln \Lambda/\mu, \lambda) + p^2 c_2(m^2/\mu^2, \ln \Lambda/\mu, \lambda) \\ \mathcal{V}_4(\Lambda; p_1, \cdots, p_4) & \longrightarrow & a_4(m^2/\mu^2, \ln \Lambda/\mu, \lambda) \\ \mathcal{V}_{2n\ge 6}(\Lambda; p_1, \cdots, p_{2n}) & \longrightarrow & 0 \end{array} \right.$$

The Λ independent part of the coefficients b_2, c_2, a_4 are not determined by the differential equation, and it must be fixed by convention. We have particularly favored the mass independent scheme in which all the coefficient functions a_2, b_2, c_2, a_4 are free of $\frac{m^2}{\mu^2}$, and

$$b_2 = c_2 = 0, \quad a_4 = -\lambda$$

at $\Lambda = \mu$.

Now, what happens if we change the convention for b_2, c_2, a_4? That would modify the action S into $S + \delta S$ which satisfies the same ERG differential equation. Now, an infinitesimal change of S is given by a composite operator

$$\delta m^2 \mathcal{O}_m + \frac{\delta z}{2} \mathcal{N} + \delta\lambda \mathcal{O}_\lambda$$

Hence, any infinitesimal change of the convention can be compensated by infinitesimal changes in m^2, λ, and the normalization of the field ϕ. Thus, we have nothing to lose by adopting a particular convention such as the mass independent scheme.

Dependence on the choice of K

In our discussions so far, we have always kept a choice of the cutoff function $K(p)$. Does physics depend on the choice of K?

Let $S_{K+\delta K}(\Lambda; m^2, \lambda; \mu)$ be the solution of the ERG equation in the mass independent scheme using the cutoff function

$$K + \delta K$$

infinitesimally different from K.
HW#7: Show that

$$S'(\Lambda) \equiv S_{K+\delta K}(\Lambda; m^2, \lambda; \mu) + \mathcal{O}_1[f] + \mathcal{O}_2[C]$$

where

$$\begin{cases} f(q) & \equiv \frac{\delta K(q/\Lambda)}{K(q/\Lambda)} \\ C(q^2) & \equiv \frac{\delta K(q/\Lambda)}{q^2+m^2} \end{cases}$$

satisfies the same ERG equation as $S_K(\Lambda; m^2, \lambda; \mu)$. Alternatively, show that

$$\begin{cases} \langle \phi(p)\phi(-p) \rangle_{S'(\Lambda)} \frac{1}{K(p/\Lambda)^2} + \frac{1-1/K(p/\Lambda)}{p^2+m^2} \\ \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_{S'(\Lambda)} \prod_{i=1}^{2n} \frac{1}{K(p_i/\Lambda)} \end{cases}$$

are Λ-independent.

This implies that the difference between $S_{K+\delta K} + \mathcal{O}_1 + \mathcal{O}_2$ and S_K is a dimension 4 composite operator, and we should be able to write

$$S_{K+\delta K}(\Lambda; m^2, \lambda; \mu) + \mathcal{O}_1[f] + \mathcal{O}_2[C] = S_K(\Lambda; m^2, \lambda; \mu) + \delta\lambda \mathcal{O}_\lambda + \delta m^2 \mathcal{O}_m + \frac{\delta z}{2} \mathcal{N}$$

in terms of some infinitesimal constants $\delta\lambda, \delta m^2$, and δz.

Let us recall the definitions of $\mathcal{O}_m, \mathcal{O}_\lambda$, and $\mathcal{N}$:

$$\begin{cases} \mathcal{O}_m & \equiv -\partial_{m^2} S - \int_q \frac{K(q/\Lambda)(1-K(q/\Lambda))}{(q^2+m^2)^2} \frac{1}{2} \left\{ \frac{\delta S}{\delta\phi(q)} \frac{\delta S}{\delta\phi(-q)} + \frac{\delta^2 S}{\delta\phi(q)\delta\phi(-q)} \right\} \\ \mathcal{O}_\lambda & \equiv -\partial_\lambda S \\ \mathcal{N} & \equiv -\int_q \phi(q) \frac{\delta S}{\delta\phi(q)} - \int_q \frac{K(q/\Lambda)(1-K(q/\Lambda))}{q^2+m^2} \left\{ \frac{\delta S}{\delta\phi(q)} \frac{\delta S}{\delta\phi(-q)} + \frac{\delta^2 S}{\delta\phi(q)\delta\phi(-q)} \right\} \end{cases}$$

Substituting this into the above, we obtain

$$S_{K+\delta K}(\Lambda; m^2 + \delta m^2, \lambda + \delta\lambda; \mu) = S_K(\Lambda; m^2, \lambda; \mu) - \mathcal{O}_1[g] - \mathcal{O}_2[D]$$

where

$$\begin{cases} g(q) & \equiv \frac{\delta K(q/\Lambda)}{K(q/\Lambda)} + \frac{1}{2}\delta z \\ D(q^2) & \equiv \frac{\delta K(q/\Lambda)}{q^2+m^2} + \frac{K(q/\Lambda)(1-K(q/\Lambda)}{q^2+m^2} \left(\delta z + \frac{\delta m^2}{q^2+m^2} \right) \end{cases}$$

Recalling

$$\begin{cases} \langle \mathcal{O}_1[g]\phi(p_1) \cdots \phi(p_{2n}) \rangle_S & = -\sum_{i=1}^{2n} g(p_i) \langle \phi(p_1) \cdots \phi(p_{2n}) \rangle_S \\ \langle \mathcal{O}_2[D]\phi(p)\phi(-p) \rangle_S & = D(p^2) \\ \langle \mathcal{O}_2[D]\phi(p_1) \cdots \phi(p_{2n}) \rangle_S & = 0 \quad (n \geq 2) \end{cases}$$

we obtain

$$
\begin{aligned}
&\langle \phi(p)\phi(-p)\rangle_{S_{K+\delta K}(\Lambda;m^2+\delta m^2,\lambda+\delta\lambda;\mu)} \\
&\quad = \langle \phi(p)\phi(-p)\rangle_{S_K(\Lambda;m^2,\lambda;\mu)} \cdot \left(1+\delta z + \frac{2\delta K(p/\Lambda)}{K(p/\Lambda)}\right) \\
&\qquad\quad - \frac{\delta K(p/\Lambda)}{p^2+m^2} - \frac{K(p/\Lambda)(1-K(p/\Lambda))}{p^2+m^2}\left(\delta z + \frac{\delta m^2}{p^2+m^2}\right) \\
&\langle \phi(p_1)\cdots\phi(p_{2n})\rangle_{S_{K+\delta K}(\Lambda;m^2+\delta m^2,\lambda+\delta\lambda;\mu)} \\
&\quad = \langle \phi(p_1)\cdots\phi(p_{2n})\rangle_{S_K(\Lambda;m^2,\lambda;\mu)} \cdot \left(1+n\delta z + \sum_{i=1}^{2n}\frac{\delta K(p_i/\Lambda)}{K(p_i/\Lambda)}\right) \qquad (n>1)
\end{aligned}
$$

The extra terms for the two-point function is just what we need to get the following results for the Λ independent correlation functions:

$$\langle \phi(p_1)\cdots\phi(p_{2n})\rangle^{K+\delta K}_{m^2+\delta m^2,\lambda+\delta\lambda;\mu} = (1+n\delta z)\,\langle \phi(p_1)\cdots\phi(p_{2n})\rangle^{K}_{m^2,\lambda;\mu} \qquad (n\geq 1)$$

where we have written $m^2, \lambda; \mu$ instead of ∞ to denote the Λ independent correlation functions. Thus, an infinitesimal change of the cutoff function K can be compensated by infinitesimal changes of m^2, λ, and the normalization of ϕ. **HW#8**: Derive the final results. (straightforward but as complicated as HW#7)

3 Application to QED

As we have discussed in subsect. 2.7, the correlation functions in the continuum limit $\Lambda \to \infty$ can be computed using the action $S(\Lambda)$ with a finite Λ. Hence, if the continuum limit has any symmetry, it must be there also in $S(\Lambda)$. As a concrete example, we study QED, the theory of photons and electrons, which has the U(1) gauge symmetry.

3.1 Perturbative construction

To construct QED, we need a real vector field A_μ for the photon and spinor fields ψ and $\bar{\psi}$ for the electron/positron. The free part of the action is given by

$$S_{\mathrm{free}} = -\frac{1}{2} \int_k A_\mu(k) A_\nu(-k) \frac{k^2 \delta_{\mu\nu} - \left(1 - \frac{1}{\xi}\right) k_\mu k_\nu}{K(k/\Lambda)} - \int_p \bar{\psi}(-p) \frac{\not{p} + im}{K(p/\Lambda)} \psi(p)$$

so that the free propagators are given by

$$\begin{cases} \langle A_\mu(k) A_\nu(-k) \rangle_{S_{\mathrm{free}}} & = \frac{K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1-\xi) \frac{k_\mu k_\nu}{k^2} \right) \\ \left\langle \psi(p) \bar{\psi}(-p) \right\rangle_{S_{\mathrm{free}}} & = \frac{K(p/\Lambda)}{\not{p} + im} = \frac{K(p/\Lambda)}{p^2 + m^2} \left(\not{p} - im \right) \end{cases}$$

ξ is the gauge fixing parameter, and m is the electron mass.

The ERG differential equation for S_{int} is obtained by generalizing that for the scalar theory as follows:

$$\begin{aligned} & -\Lambda \frac{\partial}{\partial \Lambda} S_{\mathrm{int}}(\Lambda) \\ = & \int_k \frac{\Delta(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1-\xi) \frac{k_\mu k_\nu}{k^2} \right) \frac{1}{2} \left\{ \frac{\delta S_{\mathrm{int}}}{\delta A_\mu(k)} \frac{\delta S_{\mathrm{int}}}{\delta A_\nu(-k)} + \frac{\delta^2 S_{\mathrm{int}}}{\delta A_\mu(k) \delta A_\nu(-k)} \right\} \\ & - \int_q \mathrm{Tr}\, \frac{\Delta(q/\Lambda)}{\not{q} + im} \left\{ \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} S_{\mathrm{int}} \cdot S_{\mathrm{int}} \frac{\overleftarrow{\delta}}{\delta \psi(q)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} S_{\mathrm{int}} \frac{\overleftarrow{\delta}}{\delta \psi(q)} \right\} \end{aligned}$$

where the minus sign for the second integral is due to the Fermi statistics, and the trace is for the spinor indices.

To solve the ERG differential equation, we must supply asymptotic conditions for renormalizability. For example, we can take

$$\begin{cases} \left. \frac{\delta S_{\mathrm{int}}}{\delta A_\mu(k) \delta A_\nu(-k)} \right|_0 & \to \quad \delta_{\mu\nu} \left\{ \Lambda^2 a_2(\ln \Lambda/\mu) + m^2 b_2(\ln \Lambda/\mu) + k^2 c_2(\ln \Lambda/\mu) \right\} \\ & \quad\quad + k_\mu k_\nu d_2(\ln \Lambda/\mu) \\ \left. \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_{\mathrm{int}} \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right|_0 & \to \quad \not{p}\, a_f(\ln \Lambda/\mu) + im\, b_f(\ln \Lambda/\mu) \\ \left. \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-(p+k))} \frac{\delta S_{\mathrm{int}}}{\delta A_\mu(k)} \frac{\overleftarrow{\delta}}{\delta \psi(p)} \right|_0 & \to \quad a_3(\ln \Lambda/\mu) \gamma_\mu \\ \left. \frac{\delta^4 S_{\mathrm{int}}}{\delta A_\mu(k_1) \cdots \delta A_\delta(k_4)} \right|_0 & \to \quad a_4(\ln \Lambda/\mu) \left(\delta_{\alpha\beta} \delta_{\gamma\delta} + \delta_{\alpha\gamma} \delta_{\beta\delta} + \delta_{\alpha\delta} \delta_{\beta\gamma} \right) \end{cases}$$

where $\Big|_0$ means evaluation at vanishing fields, and we choose the coefficient functions to be independent of m for mass independence.[11]

The Λ independent parts of the coefficients, except for a_2, are not fixed by the ERG differential equation, and we must fix them by convention. We can take

$$c_2(0) = 0$$

since this is a normalization condition on A_μ. Likewise we can take

$$a_f(0) = 0$$

as a normalization of ψ. By normalizing m, we can take

$$b_f(0) = 0$$

Now, in addition, we could also take $a_3(0) = e$, where e is the coupling constant, but we do not. We would rather introduce e through the Ward identities. We will show, in the remainder of this section, that $a_3(0)$ and the other coefficients $b_2(0), d_2(0), a_4(0)$ are all determined by the Ward identities.

3.2 Ward identities

Our starting point is the Ward identities in the continuum limit. For the photon two-point function, the Ward identity is

$$\frac{1}{\xi} k_\mu \left\langle A_\mu(-k) A_\nu(k) \right\rangle_\infty = \frac{k_\nu}{k^2}$$

For the higher-point functions, the identities are given by

$$\begin{aligned}&\frac{1}{\xi} k_\mu \left\langle A_\mu(-k) A_{\mu_1}(k_1) \cdots A_{\mu_M}(k_M) \psi(p_1) \cdots \psi(p_N) \bar\psi(-q_1) \cdots \bar\psi(-q_N) \right\rangle_\infty \\ &\quad = \frac{e}{k^2} \sum_{i=1}^N \left\{ \left\langle \cdots \psi(p_i - k) \cdots \right\rangle_\infty - \left\langle \cdots \bar\psi(-q_i - k) \cdots \right\rangle_\infty \right\}\end{aligned}$$

where either $M \geq 2$ or $N \geq 1$. Note that through these Ward identities, the coupling constant e is introduced into the theory.

The idea is simple. We want to transcribe the Ward identities of the Λ-independent correlation functions for the Λ-dependent correlation functions that are given directly by the cutoff action $S(\Lambda)$. Once we do this, it will not be hard to construct the corresponding invariance of the cutoff action.

---

[11] We have assumed the γ_5 invariance of the theory. Namely, under $\psi \to \gamma_5 \psi, \bar\psi \to -\bar\psi \gamma_5$, the theory is invariant if we change m to $-m$ at the same time. This γ_5 invariance excludes a term linear in Λ in the two-point fermionic vertex.

Photon two-point function

Let us first examine the Ward identity for the photon two-point function. Since

$$\langle A_\mu(-k)A_\nu(k)\rangle_\infty = \frac{1-1/K(k/\Lambda)}{k^2}\left(\delta_{\mu\nu}-(1-\xi)\frac{k_\mu k_\nu}{k^2}\right)+\langle A_\mu(-k)A_\nu(k)\rangle_S\,\frac{1}{K(k/\Lambda)^2}$$

the Ward identity gives

$$\frac{1}{\xi}k_\mu\left\langle A_\mu(-k)A_\nu(k)\right\rangle_S = K(k/\Lambda)\frac{k_\nu}{k^2}$$

Now, the equation of motion

$$\left\langle \frac{\delta S}{\delta A_\mu(k)}A_\nu(k)\right\rangle_S = -\delta_{\mu\nu}$$

gives

$$-\frac{1}{K(k/\Lambda)}\left(k^2\delta_{\mu\alpha}-\left(1-\frac{1}{\xi}\right)k_\mu k_\alpha\right)\langle A_\alpha(-k)A_\nu(k)\rangle_S+\left\langle\frac{\delta S_{\rm int}}{\delta A_\mu(k)}A_\nu(k)\right\rangle_S = -\delta_{\mu\nu}$$

Hence, multiplying by k_μ, we obtain

$$-\frac{1}{\xi}\frac{k^2}{K(k/\Lambda)}k_\mu\left\langle A_\mu(-k)A_\nu(k)\right\rangle_S + k_\mu\left\langle\frac{\delta S_{\rm int}}{\delta A_\mu(k)}A_\nu(k)\right\rangle_S = -k_\nu$$

Thus, the Ward identity gives

$$k_\mu\left\langle\frac{\delta S_{\rm int}}{\delta A_\mu(k)}A_\nu(k)\right\rangle_S = 0$$

This is equivalent to the Ward identity we started from.

Higher-point functions

We next consider the Ward identities for the higher-point functions. We first rewrite the left-hand side using the equation of motion, which gives

$$\left\langle\frac{\delta S}{\delta A_\mu(k)}A_{\mu_1}(k_1)\cdots\psi(p_1)\cdots\bar\psi(-q_1)\cdots\right\rangle_S = 0$$

Multiplying this by k_μ and substituting $S = S_{\rm free}+S_{\rm int}$, we can rewrite this as

$$\begin{aligned}&\frac{1}{\xi}\frac{k^2}{K(k/\Lambda)}k_\mu\left\langle A_\mu(-k)A_{\mu_1}(k_1)\cdots\psi(p_1)\cdots\bar\psi(-q_1)\cdots\right\rangle_S\\&= k_\mu\left\langle\frac{\delta S_{\rm int}}{\delta A_\mu(k)}A_{\mu_1}(k_1)\cdots\psi(p_1)\cdots\bar\psi(-q_1)\cdots\right\rangle_S\end{aligned}$$

Since

$$\boxed{J_\mu(-k) \equiv \frac{\delta S_{\rm int}}{\delta A_\mu(k)}}$$

is a composite operator corresponding to the charge current, we can rewrite the Ward identities as

$$\boxed{\begin{array}{c} k_\mu \left\langle J_\mu(-k) A_{\mu_1}(k_1) \cdots \psi(p_1) \cdots \bar{\psi}(-q_1) \cdots \right\rangle_\infty \\ = e \sum_{i=1}^N \left\{ \left\langle \cdots \psi(p_i - k) \cdots \right\rangle_\infty - \left\langle \cdots \bar{\psi}(-q_i - k) \cdots \right\rangle_\infty \right\} \end{array}}$$

It is easy to see that the Ward identity for the photon two-point function can be included as a special case $M = 1, N = 0$ of the above identities.

Operator equation

We wish to express the Ward identities as an operator equation

$$\boxed{k_\mu J_\mu(-k) = \Phi(-k)}$$

where we define the composite operator $\Phi(-k)$ by the right-hand side of the Ward identities as

$$\begin{aligned} &\left\langle \Phi(-k) A_{\mu_1}(k_1) \cdots \psi(p_1) \cdots \bar{\psi}(-q_1) \cdots \right\rangle_\infty \\ &\quad \equiv e \sum_{i=1}^N \left\{ \left\langle \cdots \psi(p_i - k) \cdots \right\rangle_\infty - \left\langle \cdots \bar{\psi}(-q_i - k) \cdots \right\rangle_\infty \right\} \end{aligned}$$

for any $M \geq 0$ and $N \geq 0$. $\Phi(-k)$ is a composite operator that converts

$$\left\{ \begin{array}{ccc} \psi(p_i) & \longrightarrow & e\psi(p_i - k) \\ \bar{\psi}(-q_i) & \longrightarrow & -e\bar{\psi}(-q_i - k) \end{array} \right.$$

As a preparation, we go back to the simpler scalar theory and construct an analogous operator that generates a shift of momentum:

$$\left\langle \Phi(-k) \phi(p_1) \cdots \phi(p_{2n}) \right\rangle_\infty = \sum_{i=1}^{2n} \left\langle \cdots \phi(p_i - k) \cdots \right\rangle_\infty$$

For $n > 1$, this gives

$$\left\langle \Phi(-k) \phi(p_1) \cdots \phi(p_{2n}) \right\rangle_S = \sum_{i=1}^{2n} \frac{K(p_i/\Lambda)}{K((p_i - k)/\Lambda)} \left\langle \cdots \phi(p_i - k) \cdots \right\rangle_S$$

Hence, an operator analogous to $\mathcal{O}_1[f]$ will do:

$$\Phi(-k) \overset{?}{=} - \int_q \frac{K(q/\Lambda)}{K((q-k)/\Lambda)} \phi(q - k) \frac{\delta S}{\delta \phi(q)}$$

But this is not quite correct, since for $n = 1$, we must obtain

$$\begin{aligned}
&\langle \Phi(-k)\phi(p+k)\phi(-p)\rangle_S \\
&= \frac{K((p+k)/\Lambda)}{K(p/\Lambda)} \langle \phi(p)\phi(-p)\rangle_S + \frac{K(p/\Lambda)}{K((p+k)/\Lambda)} \langle \phi(p+k)\phi(-p-k)\rangle_S \\
&\quad -K((p+k)/\Lambda)\frac{1-K(p/\Lambda)}{p^2+m^2} - K(p/\Lambda)\frac{1-K((p+k)/\Lambda)}{(p+k)^2+m^2}
\end{aligned}$$

To generate the extra terms, we need to add an operator analogous to $\mathcal{O}_2[C]$:

$$\begin{aligned}
\Phi(-k) &= -\int_q \frac{K(q/\Lambda)}{K((q-k)/\Lambda)}\phi(q-k)\frac{\delta S}{\delta\phi(q)} \\
&\quad -\int_q K((q+k)/\Lambda)\frac{1-K(q/\Lambda)}{q^2+m^2}\frac{\delta^2 S}{\delta\phi(-q)\delta\phi(q+k)}
\end{aligned}$$

HW#9: Check the final result.

Generalizing this result to QED, we define a type 1 operator

$$\mathcal{O}_1(-k) \equiv e\int_q \left[-S\frac{\overleftarrow{\delta}}{\delta\psi(q)}\frac{K(q/\Lambda)}{K((q-k)/\Lambda)}\psi(q-k) + \frac{K(q/\Lambda)}{K((q+k)/\Lambda)}\bar{\psi}(-q-k)\frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-q)}S \right]$$

and a type 2 operator

$$\mathcal{O}_2(-k) \equiv e\int_q \mathrm{Tr}\, U(-q-k,q)\left\{ \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-q)}S\cdot S\frac{\overleftarrow{\delta}}{\delta\psi(q+k)} + \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-q)}S\frac{\overleftarrow{\delta}}{\delta\psi(q+k)} \right\}$$

where

$$U(-q-k,q) \equiv K((q+k)/\Lambda)\frac{1-K(q/\Lambda)}{\not{q}+im} - \frac{1-K((q+k)/\Lambda)}{\not{q}+\not{k}+im}K(q/\Lambda)$$

We then define

$$\Phi(-k) \equiv \mathcal{O}_1(-k) + \mathcal{O}_2(-k)$$

This gives the desired relation

$$\begin{aligned}
&\left\langle \Phi(-k)A_{\mu_1}(k_1)\cdots\psi(p_1)\cdots\bar{\psi}(-q_1)\cdots\right\rangle_\infty \\
&= e\sum_{i=1}^N \left\{ \langle\cdots\psi(p_i-k)\cdots\rangle_\infty - \left\langle\cdots\bar{\psi}(-q_i-k)\cdots\right\rangle_\infty \right\}
\end{aligned}$$

HW#10: Check this.

To summarize, we have found that the Ward identities of QED can be expressed as a single operator equation as

$$k_\mu J_\mu(-k) = \Phi(-k)$$

This form of the Ward identity was first derived in [16].

3.3 Perturbative construction

We wish to show that the Ward identity can be satisfied by choosing appropriate values of the remaining four parameters: b_2, d_2, a_3, and a_4.

The asymptotic behavior of $k_\mu J_\mu$ is easily obtained from that of the action $S_{\rm int}(\Lambda)$:

$$\begin{cases} \frac{k_\mu \delta J_\mu(-k)}{\delta A_\nu(-k)}\Big|_0 = k_\nu \left(\Lambda^2 a_2(\ln\Lambda/\mu) + m^2 b_2(\ln\Lambda/\mu) + k^2(c_2+d_2)(\ln\Lambda/\mu)\right) \\ \frac{\overrightarrow{\delta}}{\delta\bar\psi(-(p+k))} k_\mu J_\mu(-k) \frac{\overleftarrow{\delta}}{\delta\psi(p)}\Big|_0 = \not k a_3(\ln\Lambda/\mu) \\ \frac{k_\mu \delta^3 J_\mu(-k)}{\delta A_\alpha(k_1)\delta A_\beta(k_2)\delta A_\gamma(k_3)}\Big|_0 = \left(k_\alpha\delta_{\beta\gamma} + k_\beta\delta_{\gamma\alpha} + k_\gamma\delta_{\alpha\beta}\right) a_4(\ln\Lambda/\mu) \end{cases}$$

On the other hand, the composite operator $\Phi(-k)$ is a dimension 4 scalar operator of momentum $-k$, vanishing at $k=0$.

Hence, its asymptotic behavior must have the same form as above[12]:

$$\begin{cases} \frac{\delta\Phi(-k)}{\delta A_\nu(-k)}\Big|_0 = k_\nu \left(\Lambda^2 \bar a_2(\ln\Lambda/\mu) + m^2 \bar b_2(\ln\Lambda/\mu) + k^2 \bar d_2(\ln\Lambda/\mu)\right) \\ \frac{\overrightarrow{\delta}}{\delta\bar\psi(-(p+k))} \Phi(-k) \frac{\overleftarrow{\delta}}{\delta\psi(p)}\Big|_0 = \not k \bar a_3(\ln\Lambda/\mu) \\ \frac{\Phi(-k)}{\delta A_\alpha(k_1)\delta A_\beta(k_2)\delta A_\gamma(k_3)}\Big|_0 = \left(k_\alpha\delta_{\beta\gamma} + k_\beta\delta_{\gamma\alpha} + k_\gamma\delta_{\alpha\beta}\right) \bar a_4(\ln\Lambda/\mu) \end{cases}$$

The two composite operators $k_\mu J_\mu$ and Φ are the same if and only if they have the same asymptotic behaviors. Hence, the Ward identity is equivalent to the following equations:

$$\begin{cases} b_2(0) &= \bar b_2(0) \\ d_2(0) &= \bar d_2(0) \\ a_3(0) &= \bar a_3(0) \\ a_4(0) &= \bar a_4(0) \end{cases}$$

where we have used the convention $c_2(0) = 0$.

Now, Φ is constructed in terms of $S(\Lambda)$. Hence, we obtain

$$\begin{cases} \frac{\delta\Phi(-k)}{\delta A_\nu(-k)}\Big|_0 = \int_q {\rm Tr}\, U(-q-k,q) \frac{\overrightarrow{\delta}}{\delta\bar\psi(-q)} \frac{\delta S}{\delta A_\nu(-k)} \frac{\overleftarrow{\delta}}{\delta\psi(q+k)}\Big|_0 \\ \frac{\overrightarrow{\delta}}{\delta\bar\psi(-(p+k))} \Phi(-k) \frac{\overleftarrow{\delta}}{\delta\psi(p)}\Big|_0 = e\left(1 - a_f(\ln\Lambda/\mu)\right)\not k \\ \qquad + \int_q \frac{\overrightarrow{\delta}}{\delta\bar\psi(-(p+k))} {\rm Tr}\left[U(-q-k,q)\frac{\overrightarrow{\delta}}{\delta\bar\psi(-q)} S \frac{\overleftarrow{\delta}}{\delta\psi(q+k)}\right] \frac{\overleftarrow{\delta}}{\delta\psi(p)}\Big|_0 \\ \frac{\delta^2\Phi(-k)}{\delta A_\alpha(k_1)\delta A_\beta(k_2)\delta A_\gamma(k_3)}\Big|_0 \\ \quad = \int_q {\rm Tr}\, U(-q-k,q) \frac{\overrightarrow{\delta}}{\delta\bar\psi(-q)} \frac{\delta^3 S}{\delta A_\alpha(k_1)\delta A_\beta(k_2)\delta A_\gamma(k_3)} \frac{\overleftarrow{\delta}}{\delta\psi(q+k)}\Big|_0 \end{cases}$$

---

[12] Here we are assuming what amounts to the CP invariance. We exclude the asymptotic behavior

$$\frac{\delta^2\Phi(-k)}{\delta A_\alpha(k_1)\delta A_\beta(k_2)}\Big|_0 \propto \epsilon_{\alpha\beta\gamma\delta} k_{1,\gamma} k_{2,\delta}$$

In chiral U(1) gauge theory this must vanish automatically. This is the anomaly cancellation condition.

We note $a_f(0) = 0$ by convention. Everything on the right-hand side has an extra loop integral. Hence, if we know S up to l-loop level, we can determine the right-hand side up to $(l+1)$-loop. Thus, by imposing the Ward identities, we can determine the coefficients $b_2(0), d_2(0), a_3(0), a_4(0)$ at $(l+1)$-loop level by knowing S up to l-loop. Therefore, we can construct the action iteratively with the help of the Ward identities.

One-loop calculations

At tree level, we obtain

$$b_2(0) = d_2(0) = a_4(0) = 0, \quad a_3(0) = e$$

We now compute $b_2(0), d_2(0)$ at one-loop. We find

$$\frac{\delta \Phi}{\delta A_\nu(-k)}\Big|_0 = e^2 \int_q \mathrm{Tr}\, \gamma_\nu U(-q-k, q)$$

Changing the integration variable as

$$q \longrightarrow \Lambda q$$

we obtain

$$\begin{aligned} \frac{\delta \Phi}{\delta A_\nu(-k)}\Big|_0 &= e^2 \Lambda^2 \int_q \mathrm{Tr}\, \gamma_\nu \cdot \left\{ K\left(q + \frac{k}{\Lambda}\right) \frac{1 - K(q)}{\not q + im/\Lambda} - \frac{1 - K\left(q + \frac{k}{\Lambda}\right)}{\not q + \not k/\Lambda + im/\Lambda} K(q) \right\} \\ &\to e^2 k_\nu \left[-2\Lambda^2 \int_q \frac{1}{q^2} \Delta(q)\left(1 - K(q)\right) + \left(m^2 + \frac{k^2}{3}\right) \int_q \frac{\Delta(q)}{(q^2)^2} \right] \end{aligned}$$

where we have expanded the integral in powers of m/Λ and k_μ/Λ. Using

$$\int_q \frac{\Delta(q) K(q)^n}{(q^2)^2} = \frac{1}{(4\pi)^2} \frac{2}{n+1}$$

(see subsect. 2.6), we obtain

$$b_2(0) = \frac{2e^2}{(4\pi)^2}, \quad d_2(0) = \frac{2e^2}{3(4\pi)^2}$$

Incidentally, the dominant photon mass term is given by

$$\Lambda^2 a_2 = -2\Lambda^2 \int_q \frac{\Delta(q)(1 - K(q))}{q^2}$$

corresponding to a large positive squared mass. Both a_2 and b_2 are non-vanishing, and the photon has mass terms in the action. But the mass terms are fixed by the gauge symmetry.

Next we consider $a_3(0)$ at one-loop:

$$\begin{aligned}
&\frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-(p+k))}\Phi(-k)\frac{\overleftarrow{\delta}}{\delta\psi(p)}\Big|_0 - e(1-a_f(\ln\Lambda/\mu))\not{k}\\
&= -e^2\int_q \gamma_\mu U(-q-k,q)\gamma_\nu\\
&\quad\times\frac{1-K((q-p)/\Lambda)}{(q-p)^2}\left(\delta_{\mu\nu}-(1-\xi)\frac{(q-p)_\mu(q-p)_\nu}{(q-p)^2}\right)\\
&\to -e^3\not{k}\int_q\frac{1}{(q^2)^2}\left\{\xi K(q)\left(1-K(q)\right)^2+\frac{3-\xi}{4}\left(1-K(q)\right)\Delta(q)\right\}
\end{aligned}$$

where we have rescaled the integration variables, and expanded the integral in powers of k_μ/Λ, p_μ/Λ. Hence,

$$a_3(0)-e=-e^3\left(\xi\int_q\frac{1}{(q^2)^2}K(q)(1-K(q))^2+\frac{3-\xi}{4(4\pi)^2}\right)$$

Finally, we compute $a_4(0)$:

$$\begin{aligned}
&\frac{\delta^3\Phi(-k)}{\delta A_\alpha(k_1)\delta A_\beta(k_2)\delta A_\gamma(k_3)}\Bigg|_0 = e^3\int_q \mathrm{Tr}\,U(-q-k,q)\\
&\times\left\{\gamma_\gamma\frac{1-K\left((q-k_3)/\Lambda\right)}{\not{q}-\not{k}_3+im}\gamma_\beta\frac{1-K\left((q-k_2-k_3)/\Lambda\right)}{\not{q}-\not{k}_2-\not{k}_3+im}\gamma_\alpha+(5\text{ permutations})\right\}\\
&\to 2e^4\left(k_\alpha\delta_{\beta\gamma}+k_\beta\delta_{\gamma\alpha}+k_\gamma\delta_{\alpha\beta}\right)\int_q\frac{\Delta(q/\Lambda)}{(q^2)^2}\left(1-K(q/\Lambda)\right)^2
\end{aligned}$$

Hence, we obtain

$$a_4(0)=\frac{4}{3}\frac{e^4}{(4\pi)^2}$$

3.4 BRST invariance

As a preparation for YM theories, we wish to formulate the Ward identity as BRST invariance. For this purpose we introduce ghost and antighost fields $c, \bar{c}$. These are free, and we modify the action by adding the free ghost kinetic term:

$$\bar{S}(\Lambda)\equiv S(\Lambda)-\int_k \bar{c}(-k)c(k)\frac{k^2}{K(k/\Lambda)}$$

Our Ward identity is given by

$$k_\mu J_\mu(-k)-\Phi(-k)=0$$

We multiply this by $c(k)$ and integrate over k to obtain

$$\int_k \epsilon c(k)\left(k_\mu J_\mu(-k)-\Phi(-k)\right)=0$$

where ϵ is an arbitrary Grassmann constant.

We first observe

$$\begin{aligned}
\int_k \epsilon c(k) k_\mu J_\mu(-k) &= \int_k \epsilon c(k) k_\mu \frac{\delta S_{\text{int}}}{\delta A_\mu(k)} \\
&= \int_k \epsilon c(k) k_\mu \left(\frac{\delta \bar{S}}{\delta A_\mu(k)} + \frac{1}{\xi} \frac{k_\mu k_\nu}{K(k/\Lambda)} A_\nu(k) \right) \\
&= \int_k \left(\epsilon k_\mu c(k) \frac{\delta \bar{S}}{\delta A_\mu(k)} - \frac{1}{\xi} \epsilon k_\mu A_\mu(k) \frac{\overrightarrow{\delta} \bar{S}}{\delta \bar{c}(-k)} \right)
\end{aligned}$$

Hence, this is the infinitesimal change of the action $\bar{S}$ under the following change of variables:

$$\begin{cases}
\delta_\epsilon A_\mu(k) &\equiv k_\mu \epsilon c(k) \\
\delta_\epsilon c(k) &\equiv 0 \\
\delta_\epsilon \bar{c}(-k) &\equiv -\frac{1}{\xi} k_\mu A_\mu(-k) \epsilon
\end{cases}$$

We next observe that Φ can be written as

$$\begin{aligned}
&\Phi(-k) = e \int_q \Bigg[\\
&S \frac{\overleftarrow{\delta}}{\delta \psi(q)} \left(- \frac{K(q/\Lambda)}{K((q-k)/\Lambda)} \psi(q-k) - K(q/\Lambda) \frac{1 - K((q-k)/\Lambda)}{\not{q} - \not{k} + im} \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-(q+k))} S \right) \\
&+ \left(\frac{K(q/\Lambda)}{K((q+k)/\Lambda)} \bar{\psi}(-q-k) + S \frac{\overleftarrow{\delta}}{\delta \psi(q+k)} \frac{1 - K((q+k)/\Lambda)}{\not{q} + \not{k} + im} K(q/\Lambda) \right) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} S \\
&+ \mathrm{Tr}\, K(q/\Lambda) \frac{1 - K((q-k)/\Lambda)}{\not{q} - \not{k} + im} \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-(q+k))} S \frac{\overleftarrow{\delta}}{\delta \psi(q)} \\
&- \mathrm{Tr}\, \frac{1 - K((q+k)/\Lambda)}{\not{q} + \not{k} + im} K(q/\Lambda) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} S \frac{\overleftarrow{\delta}}{\delta \psi(q+k)} \qquad \Bigg] \\
&= e \int_q K(q/\Lambda) \left[-S \frac{\overleftarrow{\delta}}{\delta \psi(q)} \Psi(q-k) + \bar{\Psi}(-q-k) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} S \right] \\
&\quad + e \int_q K(q/\Lambda) \mathrm{Tr} \left[\Psi(q-k) \frac{\overleftarrow{\delta}}{\delta \psi(q)} - \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} \bar{\Psi}(-q-k) \right]
\end{aligned}$$

where

$$\begin{cases}
\Psi(p) &\equiv \psi(p) + \frac{1 - K(p/\Lambda)}{\not{p} + im} \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-p)} S_{\text{int}} \\
\bar{\Psi}(-p) &\equiv \bar{\psi}(-p) + S_{\text{int}} \frac{\overleftarrow{\delta}}{\delta \psi(p)} \frac{1 - K(p/\Lambda)}{\not{p} + im}
\end{cases}$$

are the composite operators corresponding to $\psi(p), \bar{\psi}(-p)$, respectively.

HW#11: Check the derivation.

Hence, we obtain

$$\begin{aligned} - \int_k \epsilon c(k) \Phi(-k) &= \int_p \left[S \frac{\overleftarrow{\delta}}{\delta \psi(q)} \delta_\epsilon \psi(q) + \delta_\epsilon \bar{\psi}(-q) \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} S \right] \\ &\quad - \int_q \mathrm{Tr} \left[\delta_\epsilon \psi(q) \frac{\overleftarrow{\delta}}{\delta \psi(q)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} \delta_\epsilon \bar{\psi}(-q) \right] \end{aligned}$$

where we define

$$\left\{ \begin{array}{rcl} \delta_\epsilon \psi(p) & \equiv & eK(p/\Lambda) \int_k \epsilon c(k) \Psi(p-k) \\ \delta_\epsilon \bar{\psi}(-p) & \equiv & -eK(p/\Lambda) \int_k \epsilon c(k) \bar{\Psi}(-p-k) \end{array} \right.$$

Thus, altogether, the Ward identity can be written as the BRST invariance

$$\delta_\epsilon \bar{S} - \int_q \mathrm{Tr} \left[\delta_\epsilon \psi(q) \frac{\overleftarrow{\delta}}{\delta \psi(q)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} \delta_\epsilon \bar{\psi}(-q) \right] = 0$$

where $\delta_\epsilon \bar{S}$ is the infinitesimal change of the action $\bar{S}$ under the BRST transformation that is given by

$$\left\{ \begin{array}{rcl} \delta_\epsilon A_\mu(k) & = & k_\mu \epsilon c(k) \\ \delta_\epsilon c(k) & = & 0 \\ \delta_\epsilon \bar{c}(-k) & = & -\frac{1}{\xi} \epsilon k_\mu A_\mu(-k) \\ \delta_\epsilon \psi(p) & = & K(p/\Lambda) \cdot e \int_k \epsilon c(k) \Psi(p-k) \\ \delta_\epsilon \bar{\psi}(-p) & = & K(p/\Lambda) \cdot (-e) \int_k \epsilon c(k) \bar{\Psi}(-p-k) \end{array} \right.$$

Note that the second term of the BRST invariance is the jacobian of the BRST transformation. Hence, we can interpret the BRST invariance as the "quantum" invariance of $\bar{S}$ under the BRST transformation. By adopting the notation

$$\Delta_\epsilon \bar{S} \equiv - \int_q \mathrm{Tr} \left[\delta_\epsilon \psi(q) \frac{\overleftarrow{\delta}}{\delta \psi(q)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} \delta_\epsilon \bar{\psi}(-q) \right]$$

the "quantum" BRST invariance of the action is written as

$$\delta_\epsilon \bar{S} + \Delta_\epsilon \bar{S} = 0$$

The above BRST transformation of the fields has two problems:

1. it is not nilpotent (For example, $\delta_\epsilon \delta_\epsilon \psi \neq 0$.)
2. asymmetry — the transformation of $\psi, \bar{\psi}$ is given by the cutoff function times a composite operator, but that of either A_μ or $\bar{c}$ is not.

The first problem is not so serious for QED, but nilpotency is more important for YM theories; it assures the possibility of satisfying the BRST invariance by adjusting available parameters. To obtain nilpotency, we need to introduce external sources that generate BRST transformation. We will discuss it in the next subsection.

The second problem can be solved. We modify the BRST transformation of $A_\mu, \bar{c}$ as follows:

$$\left\{ \begin{array}{rcl} \delta'_\epsilon A_\mu(k) & = & K(k/\Lambda) k_\mu \epsilon c(k) \\ \delta'_\epsilon \bar{c}(-k) & = & K(k/\Lambda) \frac{-1}{\xi} \epsilon k_\mu \mathcal{A}_\mu(-k) \end{array} \right.$$

where

$$\mathcal{A}_\mu(k) \equiv A_\mu(-k) + \frac{1 - K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}(1-\xi) \right) \frac{\delta S_{\mathrm{int}}}{\delta A_\nu(-k)}$$

is the composite operator corresponding to $A_\mu(-k)$. Since $c, \bar{c}$ are non-interacting, c is already a composite operator.[13]

The action is invariant under the new transformation if it is also invariant under the difference of the two transformations:

$$\left\{ \begin{array}{rcccl} \delta''_\epsilon A_\mu(k) & \equiv & (\delta'_\epsilon - \delta_\epsilon) A_\mu(-k) & = & \left(K(k/\Lambda) - 1\right) k_\mu \epsilon c(k) \\ \delta''_\epsilon \bar{c}(-k) & \equiv & (\delta'_\epsilon - \delta_\epsilon) \bar{c}(-k) & = & \epsilon \frac{-1}{\xi} k_\mu \left(K(k/\Lambda) \mathcal{A}_\mu(-k) - A_\mu(-k) \right) \end{array} \right.$$

Since

$$k_\mu \mathcal{A}_\mu(-k) = k_\mu A_\mu(-k) + \xi \frac{1 - K(k/\Lambda)}{k^2} k_\mu \frac{\delta S_{\mathrm{int}}}{\delta A_\mu(k)}$$

we find

$$\begin{aligned} & \frac{1}{\xi} k_\mu \left(K(k/\Lambda) \mathcal{A}_\mu(-k) - A_\mu(-k) \right) \\ & = K(k/\Lambda) \left(1 - K(k/\Lambda)\right) \left(- \frac{1}{K(k/\Lambda)} \frac{1}{\xi} k_\mu A_\mu(-k) + \frac{1}{k^2} k_\mu \frac{\delta S_{\mathrm{int}}}{\delta A_\mu(k)} \right) \\ & = K(k/\Lambda) \left(1 - K(k/\Lambda)\right) \frac{1}{k^2} k_\mu \frac{\delta \bar{S}}{\delta A_\mu(k)} \end{aligned}$$

Hence, we find

$$\left\{ \begin{array}{rcl} \delta''_\epsilon A_\mu(k) & = & \left(K(k/\Lambda) - 1\right) k_\mu \epsilon c(k) \\ \delta''_\epsilon \bar{c}(-k) & = & \left(K(k/\Lambda) - 1\right) K(k/\Lambda) \epsilon \frac{1}{k^2} k_\mu \frac{\delta \bar{S}}{\delta A_\mu(k)} \end{array} \right.$$

It is easy to check the invariance of $\bar{S}$ under this:

$$\begin{aligned} \delta'' \bar{S} &= \int_k \left(\delta''_\epsilon A_\mu(k) \frac{\delta \bar{S}}{\delta A_\mu(k)} + \delta''_\epsilon \bar{c}(-k) \frac{\overrightarrow{\delta}}{\delta \bar{c}(-k)} \bar{S} \right) \\ &= \int_k \left(K(k/\Lambda) - 1\right) \left(k_\mu \epsilon c(k) \frac{\delta \bar{S}}{\delta A_\mu(k)} + K(k/\Lambda) \epsilon \frac{k_\mu}{k^2} \frac{\delta \bar{S}}{\delta A_\mu(k)} (-) \frac{k^2}{K(k/\Lambda)} c(k) \right) \\ &= 0 \end{aligned}$$

---

[13] The composite operator corresponding to c is $c + \frac{1-K}{k^2} \frac{\overrightarrow{\delta}}{\delta \bar{c}} S_{\mathrm{int}} = c$.

Thus, the action $\bar{S}$ is quantum invariant under the following new BRST transformation[14]:

$$\boxed{\begin{array}{rcl} \delta_\epsilon A_\mu(k) &=& K(k/\Lambda)\cdot k_\mu \epsilon c(k) \\ \delta_\epsilon \bar{c}(-k) &=& K(k/\Lambda)\cdot \frac{-1}{\xi}\epsilon k_\mu \mathcal{A}_\mu(-k) \\ \delta_\epsilon \psi(p) &=& K(p/\Lambda)\cdot e\int_k \epsilon c(k)\Psi(p-k) \\ \delta_\epsilon \bar{\psi}(-p) &=& K(p/\Lambda)\cdot(-e)\int_k \epsilon c(k)\bar{\Psi}(-p-k) \end{array}}$$

3.5 BRST with external sources

The most elegant way of formulating BRST invariance uses external sources that generate the BRST transformation. It is Becchi who first introduced the BRST invariance of the cutoff action in the presence of external sources.[3] Let $A^*_\mu(-k)$ be the fermionic source that generates the BRST transformation of $A_\mu(k)$. Likewise, let us introduce the bosonic sources $\bar{c}^*$, ψ^*, and $\bar{\psi}^*$. The purpose of this subsection is to show that the BRST invariance of QED can be formulated in the following form:

$$\begin{aligned} &\mathrm{e}^{-\mathcal{S}}\int_p K(p/\Lambda)\left(\frac{\overrightarrow{\delta}}{\delta A^*_\mu(-p)}\frac{\delta}{\delta A_\mu(p)} + \frac{\delta}{\delta \bar{c}^*(p)}\frac{\overrightarrow{\delta}}{\delta \bar{c}(-p)}\right. \\ &\qquad \left. + \frac{\delta}{\delta \psi^*_a(-p)}\frac{\overrightarrow{\delta}}{\delta \psi_a(p)} + \frac{\delta}{\delta \bar{\psi}^*_a(p)}\frac{\overrightarrow{\delta}}{\delta \bar{\psi}_a(-p)}\right)\mathrm{e}^{\mathcal{S}} \\ &= \int_k K(k/\Lambda)\left[\frac{\overrightarrow{\delta}}{\delta A^*_\mu(-k)}\mathcal{S}\cdot\frac{\delta\mathcal{S}}{\delta A_\mu(k)} + \frac{\delta\mathcal{S}}{\delta\bar{c}^*(k)}\frac{\overrightarrow{\delta}}{\delta\bar{c}(-k)}\mathcal{S}\right] \\ &\quad - \int_q K(q/\Lambda)\,\mathrm{Tr}\left[\frac{\delta\mathcal{S}}{\delta\psi^*(-q)}\cdot\mathcal{S}\frac{\overleftarrow{\delta}}{\delta\psi(q)} + \frac{\delta}{\delta\psi^*(-q)}\mathcal{S}\frac{\overleftarrow{\delta}}{\delta\psi(q)}\right] \\ &\quad + \int_q K(q/\Lambda)\,\mathrm{Tr}\left[\frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-p)}\mathcal{S}\cdot\frac{\delta\mathcal{S}}{\delta\bar{\psi}^*(q)} + \frac{\overrightarrow{\delta}}{\delta\bar{\psi}(-q)}\frac{\delta\mathcal{S}}{\delta\bar{\psi}^*(q)}\right] = 0 \end{aligned}$$

where $\mathcal{S}$ is the action with external sources. The meaning of this equation is clear. The BRST transformation of the fields is given by functional derivatives with respect to external sources:

$$\left\{\begin{array}{rcl} \delta_\epsilon A_\mu(k) &=& \epsilon K(k/\Lambda)\frac{\overrightarrow{\delta}}{\delta A^*_\mu(-k)}\mathcal{S} \\ \delta_\epsilon \bar{c}(-k) &=& \epsilon K(k/\Lambda)\frac{\delta}{\delta \bar{c}^*(k)}\mathcal{S} \\ \delta_\epsilon \psi(q) &=& \epsilon K(q/\Lambda)\frac{\delta}{\delta \psi^*(-q)}\mathcal{S} \\ \delta_\epsilon \bar{\psi}(-q) &=& \epsilon K(q/\Lambda)\frac{\delta}{\delta \bar{\psi}^*(q)}\mathcal{S} \end{array}\right.$$

---

[14] The jacobian of $\delta_\epsilon A_\mu, \delta_\epsilon \bar{c}$ is 1.

At the vanishing sources, we must find

$$\left\{ \begin{array}{rcl} \frac{\overrightarrow{\delta}}{\delta A_\mu^*(-k)}\mathcal{S} &=& k_\mu c(k) \\ \frac{\delta}{\delta \bar{c}^*(k)}\mathcal{S} &=& -\frac{1}{\xi} k_\mu \mathcal{A}_\mu(-k) \\ \frac{\delta}{\delta \psi^*(-q)}\mathcal{S} &=& e \int_k c(k)\Psi(q-k) \\ \frac{\delta}{\delta \bar{\psi}^*(q)}\mathcal{S} &=& -e \int_k c(k)\bar{\Psi}(-q-k) \end{array} \right.$$

In this subsection we wish to understand the general structure of the BRST transformation generated by sources; especially we would like to introduce a BRST transformation δ_Q that is nilpotent. In the next subsection, we will construct $\mathcal{S}$ explicitly in terms of $\bar{S}$ without sources that we have already constructed.

In order to understand the main properties of the system, it is best to treat the simplest possible system with the same structure. So, instead of QED, we consider the theory of a scalar field ϕ in the presence of an external fermionic source ϕ^*. Let $\mathcal{S}[\phi, \phi^*]$ be a cutoff action satisfying the Polchinski ERG differential equation

$$\begin{aligned} -\Lambda \frac{\partial}{\partial \Lambda}\mathcal{S} = \int_q \frac{\Delta(q/\Lambda)}{q^2+m^2} \Bigg(& \frac{q^2+m^2}{K(q/\Lambda)} \phi(q) \frac{\delta \mathcal{S}}{\delta \phi(-q)} \\ & + \frac{1}{2} \left\{ \frac{\delta \mathcal{S}}{\delta \phi(q)} \frac{\delta \mathcal{S}}{\delta \phi(-q)} + \frac{\delta^2 \mathcal{S}}{\delta \phi(q) \delta \phi(-q)} \right\} \Bigg) \end{aligned}$$

Note that the external source is a purely classical source that does not participate in the ERG equation. Differentiating the equation with respect to ϕ^*, we find that $\frac{\overrightarrow{\delta}}{\delta \phi^*(-p)}\mathcal{S}$ is a composite operator. We also recall that $K(p/\Lambda)\frac{\delta \mathcal{S}}{\delta \phi(-p)}$ is a composite operator.

Now, as a preparation, we wish to show

1. If $\mathcal{O}$ is a composite operator, so is

$$A \equiv K(q/\Lambda) \left(\mathcal{O} \frac{\delta \mathcal{S}}{\delta \phi(-q)} + \frac{\delta \mathcal{O}}{\delta \phi(-q)} \right) = \mathrm{e}^{-\mathcal{S}} K(q/\Lambda) \frac{\delta}{\delta \phi(-q)} \left(\mathcal{O} \mathrm{e}^{\mathcal{S}} \right)$$

2. If $\mathcal{O}$ is a composite operator, so is

$$B \equiv \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \mathcal{O} + \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \mathcal{S} \cdot \mathcal{O} = \mathrm{e}^{-\mathcal{S}} \frac{\overrightarrow{\delta}}{\delta \phi^*(-p)} \left(\mathcal{O} \mathrm{e}^{\mathcal{S}} \right)$$

(Proof) Let us consider the Λ derivative of A:

$$\begin{aligned} & -\Lambda \frac{\partial}{\partial \Lambda} \left(K(q/\Lambda) \left(\mathcal{O} \frac{\delta \mathcal{S}}{\delta \phi(-q)} + \frac{\delta \mathcal{O}}{\delta \phi(-q)} \right) \right) \\ & = \mathcal{D}\mathcal{O} \cdot K(q/\Lambda) \frac{\delta \mathcal{S}}{\delta \phi(-q)} + \mathcal{O} \cdot K(q/\Lambda) \mathcal{D} \frac{\delta \mathcal{S}}{\delta \phi(-q)} \end{aligned}$$

$$
\begin{aligned}
&+K(q/\Lambda)\frac{\delta}{\delta\phi(-q)}\mathcal{D}\mathcal{O}-\Delta(q/\Lambda)\frac{\delta}{\delta\phi(-q)}\mathcal{O}\\
&=\mathcal{D}\left(\mathcal{O}K(q/\Lambda)\frac{\delta\mathcal{S}}{\delta\phi(-q)}\right)-\int_r\frac{\Delta(r/\Lambda)}{r^2+m^2}\frac{\delta}{\delta\phi(r)}\mathcal{O}\cdot K(q/\Lambda)\frac{\delta^2\mathcal{S}}{\delta\phi(-r)\delta\phi(-q)}\\
&\quad+K(q/\Lambda)\frac{\delta}{\delta\phi(-q)}\mathcal{D}\mathcal{O}-\Delta(q/\Lambda)\frac{\delta}{\delta\phi(-q)}\mathcal{O}\\
&=\mathcal{D}\left(\mathcal{O}K(q/\Lambda)\frac{\delta\mathcal{S}}{\delta\phi(-q)}\right)-\int_r\frac{\Delta(r/\Lambda)}{r^2+m^2}\frac{\delta}{\delta\phi(r)}\mathcal{O}\cdot K(q/\Lambda)\frac{\delta^2\mathcal{S}_{\mathrm{int}}}{\delta\phi(-r)\delta\phi(-q)}\\
&\quad+K(q/\Lambda)\frac{\delta}{\delta\phi(-q)}\mathcal{D}\mathcal{O}
\end{aligned}
$$

Since

$$
\frac{\delta}{\delta\phi(-q)}\mathcal{D}\mathcal{O}=\mathcal{D}\frac{\delta}{\delta\phi(-q)}\mathcal{O}+\int_r\frac{\Delta(r/\Lambda)}{r^2+m^2}\frac{\delta^2\mathcal{S}_{\mathrm{int}}}{\delta\phi(-r)\delta\phi(-q)}\frac{\delta}{\delta\phi(r)}\mathcal{O}
$$

we obtain

$$
-\Lambda\frac{\partial}{\partial\Lambda}A=\mathcal{D}A
$$

Hence, A is a composite operator.

We next consider B:

$$
\begin{aligned}
&-\Lambda\frac{\partial}{\partial\Lambda}\left(\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\mathcal{O}+\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\mathcal{S}\cdot\mathcal{O}\right)\\
&=\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\left(\mathcal{D}\mathcal{O}\right)+\mathcal{D}\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\mathcal{S}\cdot\mathcal{O}+\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\mathcal{S}\cdot\mathcal{D}\mathcal{O}\\
&=\mathcal{D}\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\mathcal{O}+\int_q\frac{\Delta(q/\Lambda)}{q^2+m^2}\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\frac{\delta\mathcal{S}_{\mathrm{int}}}{\delta\phi(-q)}\cdot\frac{\delta}{\delta\phi(q)}\mathcal{O}\\
&\quad+\mathcal{D}\left(\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\mathcal{S}\cdot\mathcal{O}\right)-\int_q\frac{\Delta(q/\Lambda)}{q^2+m^2}\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\frac{\delta\mathcal{S}_{\mathrm{int}}}{\delta\phi(-q)}\cdot\frac{\delta}{\delta\phi(q)}\mathcal{O}\\
&=\mathcal{D}B
\end{aligned}
$$

Hence, B is a composite operator. **Q.E.D.**

HW#12: Alternatively you can show

$$
\begin{aligned}
\langle A(q)\phi(p_1)\cdots\phi(p_{2n})\rangle_\infty&=-\sum_{i=1}^{2n}(2\pi)^4\delta^{(4)}(p_i+q)\left\langle\cdots\widehat{\phi(p_i)}\mathcal{O}\cdots\right\rangle_\infty\\
\langle B(p)\phi(p_1)\cdots\phi(p_{2n})\rangle_\infty&=\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)}\langle\mathcal{O}\,\phi(p_1)\cdots\phi(p_{2n})\rangle_\infty
\end{aligned}
$$

This shows A and B are composite.

Nilpotent δ_Q

Now, let us apply what we have found. Given a composite operator $\mathcal{O}$,

$$X(p) \equiv K(p/\Lambda) \mathrm{e}^{-\mathcal{S}} \frac{\delta}{\delta\phi(-p)} \left(\mathrm{e}^{\mathcal{S}} \mathcal{O}\right) = K(p/\Lambda) \left(\frac{\delta \mathcal{S}}{\delta\phi(-p)} \mathcal{O} + \frac{\delta \mathcal{O}}{\delta\phi(-p)} \right)$$

is also a composite operator according to 1. Applying 2 to X, we obtain

$$Y(p) \equiv \mathrm{e}^{-\mathcal{S}} \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \left(\mathrm{e}^{\mathcal{S}} X\right) = \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} X(p) + \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \mathcal{S} \cdot X(p)$$

is also a composite operator. We find

$$\begin{aligned} Y(p) &= K(p/\Lambda) \mathrm{e}^{-\mathcal{S}} \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \frac{\delta}{\delta\phi(-p)} \left(\mathrm{e}^{\mathcal{S}} \mathcal{O}\right) \\ &= K(p/\Lambda) \Bigg[\left(\frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \mathcal{S} \cdot \frac{\delta \mathcal{S}}{\delta\phi(-p)} + \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \frac{\delta \mathcal{S}}{\delta\phi(-p)} \right) \mathcal{O} \\ &\qquad + \frac{\delta \mathcal{S}}{\delta\phi(-p)} \cdot \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \mathcal{O} + \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \mathcal{S} \cdot \frac{\delta \mathcal{O}}{\delta\phi(-p)} + \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \frac{\delta \mathcal{O}}{\delta\phi(-p)} \Bigg] \end{aligned}$$

Since

$$K(p/\Lambda) \frac{\delta \mathcal{S}}{\delta\phi(-p)}$$

is a composite operator, we find that

$$K(p/\Lambda) \left(\frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \frac{\delta \mathcal{S}}{\delta\phi(-p)} + \frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \mathcal{S} \cdot \frac{\delta \mathcal{S}}{\delta\phi(-p)} \right)$$

is also a composite operator, according to 2. (Alternatively, apply 1 to the composite operator $\frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \mathcal{S}$.) Thus, it is consistent with ERG to assume that its integral over p vanishes:

$$\Sigma[\phi, \phi^*] \equiv \int_p K(p/\Lambda) \left(\frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \frac{\delta S}{\delta\phi(-p)} + \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} S \cdot \frac{\delta S}{\delta\phi(-p)} \right) = 0$$

This is BRST invariance. If this equation is satisfied at a particular Λ, the ERG differential equation guarantees that it is satisfied at an arbitrary Λ. Becchi's BRST invariance has precisely this form.

Assuming that the above BRST invariance is satisfied, we define

$$\begin{aligned} &\delta_Q \mathcal{O} \equiv \int_p Y(p) \\ &= \int_p K(p/\Lambda) \left[\frac{\delta \mathcal{S}}{\delta\phi(-p)} \cdot \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \mathcal{O} + \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \mathcal{S} \cdot \frac{\delta \mathcal{O}}{\delta\phi(-p)} + \frac{\overrightarrow{\delta}}{\delta\phi^*(p)} \frac{\delta \mathcal{O}}{\delta\phi(-p)} \right] \end{aligned}$$

Hence, given a composite operator $\mathcal{O}$, $\delta_Q \mathcal{O}$ defines a composite operator. We define δ_Q by the above, even for $\mathcal{O}$ which is not a composite operator. It is straightforward to show that δ_Q is nilpotent:

$$\boxed{\delta_Q \delta_Q = 0} \qquad \text{assuming } \Sigma = 0$$

as long as the BRST invariance holds.
HW#13: Show the nilpotency. ($\Sigma = 0$ is necessary.)

Suppose $\Sigma \neq 0$ so that δ_Q is not necessarily nilpotent. We still get an algebraic constraint on Σ:

$$\boxed{\delta_Q \Sigma[\phi, \phi^*] = 0} \qquad \text{without assuming } \Sigma = 0$$

HW#14: Derive this.

Now, Becchi's BRST invariance can be interpreted as the invariance (including the functional measure) of $\mathcal{S}$ under the infinitesimal field transformation:

$$\delta\phi(p) \equiv \delta_Q \phi(p) = K(p/\Lambda) \cdot \underbrace{\frac{\overrightarrow{\delta}}{\delta\phi^*(-p)} \mathcal{S}}_{\text{composite operator}}$$

The nilpotency $\delta_Q^2 = 0$, guaranteed by the BRST invariance, implies

$$\delta_Q (\delta\phi(p)) = 0$$

HW#15: Calculate this explicitly and show that $\Sigma = 0$ is necessary for the equality.

Let us now consider generalizing the above results for a theory with many fields ϕ_i either bosonic or fermionic. Let ϕ_i^* be the corresponding source with the opposite statistics. The BRST invariance of the action is given by

$$\Sigma[\phi, \phi^*] \equiv \mathrm{e}^{-\mathcal{S}} \int_q \sum_i \frac{\overrightarrow{\delta}}{\delta\phi_i^*(-q)} \frac{\overrightarrow{\delta}}{\delta\phi_i(q)} \mathrm{e}^{\mathcal{S}} = 0$$

Since

$$Y(p) = \mathrm{e}^{-\mathcal{S}} K(p/\Lambda) \sum_i \frac{\overrightarrow{\delta}}{\delta\phi_i^*(-p)} \frac{\overrightarrow{\delta}}{\delta\phi_i(p)} \left(\mathrm{e}^{\mathcal{S}} \mathcal{O} \right)$$

we arrive at the definition of δ_Q given by

$$\begin{aligned} \delta_Q \mathcal{O} \quad \equiv \quad & \int_p K(p/\Lambda) \Bigg(\frac{\overrightarrow{\delta}}{\delta\phi_i^*(-p)} \mathcal{S} \cdot \frac{\overrightarrow{\delta}}{\delta\phi_i(p)} \mathcal{O} + \frac{\overrightarrow{\delta}}{\delta\phi_i(p)} \mathcal{S} \cdot \frac{\overrightarrow{\delta}}{\delta\phi_i^*(-p)} \mathcal{O} \\ & + \frac{\overrightarrow{\delta}}{\delta\phi_i^*(-p)} \frac{\overrightarrow{\delta}}{\delta\phi_i(p)} \mathcal{O} \Bigg) \end{aligned}$$

Especially, we obtain

$$\delta_Q \phi_i(p) = K(p/\Lambda) \frac{\overrightarrow{\delta}}{\delta \phi_i^*(-p)} \mathcal{S}$$

so that the BRST invariance is written as the quantum invariance of the action under the BRST transformation:

$$\Sigma[\phi, \phi^*] \equiv \int_q \left(\delta_Q \phi_i(q) \cdot \frac{\overrightarrow{\delta}}{\delta \phi_i(q)} \mathcal{S} + \frac{\overrightarrow{\delta}}{\delta \phi_i(q)} \delta_Q \phi_i(q) \right) = 0$$

Finally, in the case of QED, the BRST invariance has been already given at the beginning of this subsection. We have used

$$\frac{\overrightarrow{\delta}}{\delta A_\mu^*(-k)} \frac{\delta \mathcal{S}}{\delta A_\mu(k)} = 0, \quad \frac{\delta}{\delta \bar{c}^*(k)} \frac{\overrightarrow{\delta}}{\delta \bar{c}(-k)} \mathcal{S} = 0$$

which we will verify in the next subsection. δ_Q is defined as follows:

$$\begin{aligned}
&\delta_Q \mathcal{O} \equiv \int_q K(q/\Lambda) \Bigg[\\
&\frac{\delta \mathcal{S}}{\delta A_\mu(-q)} \cdot \frac{\overrightarrow{\delta}}{\delta A_\mu^*(q)} \mathcal{O} + \frac{\overrightarrow{\delta}}{\delta A_\mu^*(q)} \mathcal{S} \cdot \frac{\delta}{\delta A_\mu(-q)} \mathcal{O} + \frac{\overrightarrow{\delta}}{\delta A_\mu^*(q)} \frac{\delta \mathcal{O}}{\delta A_\mu(-q)} \\
&+ \frac{\overrightarrow{\delta}}{\delta \bar{c}(-q)} \mathcal{S} \cdot \frac{\delta}{\delta \bar{c}^*(q)} \mathcal{O} + \frac{\delta \mathcal{S}}{\delta \bar{c}^*(q)} \frac{\overrightarrow{\delta}}{\delta \bar{c}(-q)} \mathcal{O} + \frac{\delta}{\delta \bar{c}^*(k)} \frac{\overrightarrow{\delta}}{\delta \bar{c}(-k)} \mathcal{O} \\
&-(-)^F \mathrm{Tr} \left(\frac{\delta \mathcal{O}}{\delta \psi^*(-q)} \mathcal{S} \frac{\overleftarrow{\delta}}{\delta \psi(q)} + \mathcal{O} \frac{\overleftarrow{\delta}}{\delta \psi(q)} \frac{\delta \mathcal{S}}{\delta \psi^*(-q)} + \frac{\delta \mathcal{O}}{\delta \psi^*(-q)} \frac{\overleftarrow{\delta}}{\delta \psi(q)} \right) \\
&+ \mathrm{Tr} \left(\frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} \mathcal{S} \cdot \mathcal{O} \frac{\overleftarrow{\delta}}{\delta \bar{\psi}^*(q)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} \mathcal{O} \cdot \mathcal{S} \frac{\overleftarrow{\delta}}{\delta \bar{\psi}^*(q)} + \frac{\overrightarrow{\delta}}{\delta \bar{\psi}(-q)} \mathcal{O} \frac{\overleftarrow{\delta}}{\delta \bar{\psi}^*(q)} \right) \Bigg]
\end{aligned}$$

where $(-)^F = \pm 1$ depending on whether $\mathcal{O}$ is bosonic or fermionic. δ_Q is nilpotent if $\mathcal{S}$ is BRST invariant.

3.6 Explicit construction of $\mathcal{S}$ in terms of $\bar{S}$

The purpose of this subsection is to construct the BRST invariant action $\mathcal{S}$ in the presence of external sources explicitly in terms of the BRST invariant action $\bar{S}$ without sources. Since we must find

$$\left\{ \begin{array}{rcl}
\frac{\overrightarrow{\delta}}{\delta A_\mu^*(-k)} \mathcal{S} &=& k_\mu c(k) \\
\frac{\delta}{\delta \bar{c}^*(k)} \mathcal{S} &=& -\frac{1}{\xi} k_\mu \mathcal{A}_\mu(-k) \\
\frac{\delta}{\delta \psi^*(-q)} \mathcal{S} &=& e \int_k c(k) \Psi(q-k) \\
\frac{\delta}{\delta \bar{\psi}^*(q)} \mathcal{S} &=& -e \int_k c(k) \bar{\Psi}(-q-k)
\end{array} \right.$$

at the vanishing sources, we obtain

$$\begin{aligned}\mathcal{S}-\bar{S} \quad \simeq \quad & \int_k \left[A_\mu^*(-k)k_\mu c(k)+\bar{c}^*(k)\frac{-1}{\xi}k_\mu \mathcal{A}_\mu(-k)\right] \\ & +e\int_{q,k}\left[\psi^*(-q)c(k)\Psi(q-k)-c(k)\bar{\Psi}(-q-k)\bar{\psi}^*(q)\right]\end{aligned}$$

to first order in external sources. The question is how to add terms higher order in external sources so that $\mathcal{S}$ satisfies both ERG and BRST invariance.

As a preparation we consider a simpler problem. Suppose we have an action of a scalar field ϕ that satisfies the ERG differential equation:

$$-\Lambda\frac{\partial S}{\partial\Lambda}=\int_q\frac{\Delta(q/\Lambda)}{q^2+m^2}\left[\frac{q^2+m^2}{K(q/\Lambda)}\phi(q)\frac{\delta S}{\delta\phi(q)}+\frac{1}{2}\left\{\frac{\delta S}{\delta\phi(q)}\frac{\delta S}{\delta\phi(-q)}+\frac{\delta^2 S}{\delta\phi(q)\delta\phi(-q)}\right\}\right]$$

We wish to introduce an external source $J(-q)$ that couples to the scalar field $\phi(q)$. Since $\frac{\delta S}{\delta J(-q)}$ at $J=0$ is a composite operator with respect to S, we find that the action $S[\phi,J]$ in the presence of the source is given by

$$S[\phi,J]-\underbrace{S[\phi,0]}_{=S[\phi]}\simeq\int_q J(-q)\Phi(q)$$

to first order in J, where Φ is the composite operator

$$\Phi(q)\equiv\phi(q)+\frac{1-K(q/\Lambda)}{q^2+m^2}\frac{\delta S_{\mathrm{int}}}{\delta\phi(-q)}$$

in the absence of J. It is not hard to guess the full source dependence of $S[\phi,J]$:

$$\begin{aligned}S[\phi,J] \quad = \quad & S_{\mathrm{free}}[\phi]+\int_q J(-q)\phi(q)+\frac{1}{2}\int_q J(-q)\frac{1-K(q/\Lambda)}{q^2+m^2}J(q) \\ & +S_{\mathrm{int}}\left[\phi(q)\to\phi(q)+J(q)\frac{1-K(q/\Lambda)}{q^2+m^2}\right]\end{aligned}$$

To first order in J, this reproduces $S[\phi,J]$ given above.

Let us verify that $S[\phi,J]$ satisfies the ERG differential equation. We obtain

$$\begin{aligned}&-\Lambda\frac{\partial}{\partial\Lambda}\left(S[\phi,J]-S_{\mathrm{free}}[\phi]\right)\\ &=\frac{1}{2}\int_q J(-q)\frac{\Delta(q/\Lambda)}{q^2+m^2}J(q)+\int_q J(q)\frac{\Delta(q/\Lambda)}{q^2+m^2}\frac{\delta S_{\mathrm{int}}}{\delta\phi(q)}\\ &\quad+\int_q\frac{\Delta(q/\Lambda)}{q^2+m^2}\frac{1}{2}\left\{\frac{\delta S_{\mathrm{int}}}{\delta\phi(q)}\frac{\delta S_{\mathrm{int}}}{\delta\phi(-q)}+\frac{\delta^2 S_{\mathrm{int}}}{\delta\phi(q)\delta\phi(-q)}\right\}\end{aligned}$$

Since

$$\left\{\begin{array}{rcl}\frac{\delta}{\delta\phi(q)}\left(S[\phi,J]-S_{\mathrm{free}}[\phi]\right) & = & J(-q)+\frac{\delta S_{\mathrm{int}}}{\delta\phi(q)}\\ \frac{\delta^2}{\delta\phi(q)\delta\phi(-q)}\left(S[\phi,J]-S_{\mathrm{free}}[\phi]\right) & = & \frac{\delta^2 S_{\mathrm{int}}}{\delta\phi(q)\delta\phi(-q)}\end{array}\right.$$

we obtain

$$-\Lambda \frac{\partial S_{\mathrm{int}}[\phi, J]}{\partial \Lambda} = \int_q \frac{\Delta(q/\Lambda)}{q^2+m^2} \frac{1}{2} \left\{ \frac{\delta S_{\mathrm{int}}[\phi, J]}{\delta \phi(q)} \frac{\delta S_{\mathrm{int}}[\phi, J]}{\delta \phi(-q)} + \frac{\delta^2 S_{\mathrm{int}}[\phi, J]}{\delta \phi(q) \delta \phi(-q)} \right\}$$

where

$$S_{\mathrm{int}}[\phi, J] \equiv S[\phi, J] - S_{\mathrm{free}}[\phi]$$

Thus, $S[\phi, J]$ satisfies the ERG differential equation.

The external source $J(q)$ is a dimension 3 parameter of the theory, which happens to be momentum dependent. Hence, J enters into the asymptotic behavior of the action only for the one-point vertex:

$$\frac{\delta S_{\mathrm{int}}[\phi, J]}{\delta \phi(-q)}\Big|_0 \to J(q)$$

Thus, the ERG differential equation determines the J-dependence of the action unambiguously. In terms of Feynman graphs, $\int_q J(-q)\phi(q)$ gives a one-point vertex



This vertex can be connected to another vertex only through a high-momentum propagator $(1-K(q/\Lambda))/(q^2+m^2)$. Multiple one-point vertices can be connected to an original vertex. Hence, a typical vertex in the presence of sources looks like the left figure below.





The only exception is two one-point vertices connected to each other by a high-momentum propagator. (the right figure above)

Now, let us apply the above result to QED. We first tabulate the fields and their external sources:

ext. source	coupled field	dimensions of the source
$k_\mu A^*_\mu(-k)$	$c(k)$	3
$-\frac{1}{\xi} k_\mu \bar{c}^*(k)$	$A_\mu(-k)$	3
$e \int_k \psi^*(-q-k) c(k)$	$\psi(q)$	5/2
$e \int_k c(k) \bar{\psi}^*(q-k)$	$\bar{\psi}(-q)$	5/2

where we treat the free field c as an external field. Then, the full action that satisfies the ERG differential equation in the presence of the external sources is given by

$$\begin{aligned}
\mathcal{S} &= \bar{S}_{\mathrm{free}} + \int_k \left[k_\mu A_\mu^*(-k) c(k) + \frac{-1}{\xi} k_\mu \bar{c}^*(k) A_\mu(-k) \right] \\
&\quad + e \int_{q,k} \left[\psi^*(-q-k) c(k) \psi(q) + \bar{\psi}(-q) c(k) \bar{\psi}^*(q-k) \right] \\
&\quad - \frac{1}{2\xi} \int_k \left(1 - K(k/\Lambda)\right) \bar{c}^*(-k) \bar{c}^*(k) \\
&\quad + e^2 \int_{q,k,l} \psi^*(-q-k) c(k) \frac{1 - K(q/\Lambda)}{\not{q} + im} c(l) \bar{\psi}^*(q-l) \\
&\quad + S_{\mathrm{int}} \Big[A_\mu(k) \to A_\mu(k) - \frac{1 - K(k/\Lambda)}{k^2} k_\mu \bar{c}^*(k), \\
&\qquad \psi(q) \to \psi(q) + \frac{1 - K(q/\Lambda)}{\not{q} + im} e \int_k c(k) \bar{\psi}^*(q-k), \\
&\qquad \bar{\psi}(-q) \to \bar{\psi}(-q) + e \int_k \psi^*(-q-k) c(k) \frac{1 - K(q/\Lambda)}{\not{q} + im} \Big]
\end{aligned}$$

We note

$$\frac{\overrightarrow{\delta}}{\delta A_\mu^*(-k)} \mathcal{S} = k_\mu c(k), \quad \frac{\overrightarrow{\delta}}{\delta \bar{c}(-k)} \mathcal{S} = -k^2 c(k)$$

Hence,

$$\int_k K(k/\Lambda) \frac{\overrightarrow{\delta}}{\delta A_\mu^*(-k)} \frac{\delta \mathcal{S}}{\delta A_\mu(k)} = \int_k K(k/\Lambda) \frac{\delta}{\delta \bar{c}^*(k)} \frac{\overrightarrow{\delta}}{\delta \bar{c}(-k)} \mathcal{S} = 0$$

The antifield dependence of the action for QED was first derived in [17], but their result corresponds to a different choice of the cutoff dependence of the external sources. The above expression has been obtained independently in [18], where its equivalence with the action of [17] has also been shown.

HW#16: Verify the BRST invariance $\Sigma = 0$ explicitly using the above action. (straightforward but tedious)

4 Application to YM theories

We now consider YM theories. We first set our notations straight. We denote the gauge fields as A^a_μ and the Faddeev-Popov ghosts by $c^a, \bar{c}^a$. We keep the gauge group general. In terms of the totally antisymmetric structure constant f^{abc}, the covariant derivatives are given by

$$\left\{ \begin{array}{rcl} F^a_{\mu\nu} & \equiv & \partial_\mu A^a_\nu - \partial_\nu A^a_\mu - g f^{abc} A^b_\mu A^c_\nu \\ (D_\mu c)^a & \equiv & \partial_\mu c^a - g f^{abc} A^b_\mu c^c \end{array} \right.$$

Hence, for the Fourier transforms, we obtain

$$\left\{ \begin{array}{rcl} F^a_{\mu\nu}(k) & = & i k_\mu A^a_\nu(k) - i k_\nu A^a_\mu(k) - g f^{abc} \int_l A^b_\mu(k-l) A^c_\nu(l) \\ (D_\mu c)^a(k) & = & i k_\mu c^a(k) - g f^{abc} \int_l A^b_\mu(k-l) c^c(l) \end{array} \right.$$

The classical action

$$\begin{aligned} \mathcal{S}_{\mathrm{classical}} \equiv & \int d^4x \left[-\frac{1}{4} F^a_{\mu\nu} F^a_{\mu\nu} - \frac{1}{2\xi} \left(\partial_\mu A^a_\mu\right)^2 - \partial_\mu \bar{c}^a \left(D_\mu c\right)^a \right. \\ & \left. + A^{a*}_\mu \frac{1}{i} \left(D_\mu c\right)^a + \bar{c}^{a*} \frac{1}{\xi} \frac{1}{i} \partial_\mu A^a_\mu + c^{a*} \frac{g}{2i} f^{abc} c^b c^c \right] \end{aligned}$$

has the BRST invariance for which the cutoff function is taken as 1. The classical BRST transformation is given by

$$\left\{ \begin{array}{rcccl} \delta_\epsilon A^a_\mu & = & \epsilon \frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu} \mathcal{S}_{\mathrm{classical}} & = & \epsilon \frac{1}{i} \left(D_\mu c\right)^a \\ \delta_\epsilon \bar{c}^a & = & \epsilon \frac{\delta}{\delta \bar{c}^{a*}} \mathcal{S}_{\mathrm{classical}} & = & \frac{1}{\xi} \epsilon \frac{1}{i} \partial \cdot A^a \\ \delta_\epsilon c^a & = & \epsilon \frac{\delta}{\delta c^{a*}} \mathcal{S}_{\mathrm{classical}} & = & \epsilon \frac{1}{2i} g f^{abc} c^b c^c \end{array} \right.$$

4.1 YM theories without sources

We first summarize the characteristic properties of the YM theories in the absence of sources. The correlation functions in the continuum limit satisfy two important relations:

1. **Ward identities**

$$\begin{aligned} & \frac{1}{\xi} k_\mu \left\langle A^a_\mu(-k) A^{a_1}_{\nu_1}(k_1) \cdots A^{a_n}_{\nu_n}(k_n) \right\rangle_\infty \\ & = \sum_{i=1}^n \left\langle A^{a_1}_{\nu_1}(k_1) \cdots \frac{1}{i} \left(D_{\nu_i} c\right)^{a_i}(k_i) \cdots A^{a_n}_{\nu_n}(k_n) \bar{c}^a(-k) \right\rangle_\infty \end{aligned}$$

   where $D_\mu c$ is the composite operator for the covariant derivative.
   (naïve proof) We apply the BRST transformation to

$$\left\langle A^{a_1}_{\nu_1}(k_1) \cdots A^{a_n}_{\nu_n}(k_n) \bar{c}^a(-k) \right\rangle_\infty$$

Since the action is invariant, we obtain

$$\begin{aligned}&\sum_{i=1}^{n}\left\langle A^{a_1}_{\nu_1}(k_1)\cdots\frac{1}{i}(D_{\nu_i}c)^{a_i}(k_i)\cdots\bar{c}^a(-k)\right\rangle_\infty\\&+\left\langle A^{a_1}_{\nu_1}(k_1)\cdots A^{a_n}_{\nu_n}(k_n)\frac{1}{\xi}(-k_\mu)A^a_\mu(-k)\right\rangle_\infty=0\end{aligned}$$

2. **The ghost equations of motion** — The covariant derivative $D_\mu c$ must satisfy

$$\begin{cases}k_\mu\left\langle\frac{1}{i}\left(D_\mu c\right)^a(k)\bar{c}^b(-k)\right\rangle_\infty=\delta^{ab}\\ k_\mu\left\langle\frac{1}{i}\left(D_\mu c\right)^a(k)A^{a_1}_{\nu_1}(k_1)\cdots A^{a_{n-1}}_{\nu_{n-1}}(k_{n-1})\bar{c}^{a_n}(k_n)\right\rangle_\infty=0\quad(n>1)\end{cases}$$

(naïve proof) Since

$$\frac{\overrightarrow{\delta}}{\delta\bar{c}^a(-k)}S=-k_\mu\frac{1}{i}(D_\mu c)^a(k)$$

we obtain the above equations of motion.

4.2 YM theories with sources

The cutoff action is given in the familiar form:

$$\mathcal{S}=S_{\text{free}}+\mathcal{S}_{\text{int}}$$

where the free action is given by

$$S_{\text{free}}=-\int_k\left[\frac{1}{2}A^a_\mu(k)A^a_\nu(-k)\frac{k^2\delta_{\mu\nu}-\left(1-\frac{1}{\xi}\right)k_\mu k_\nu}{K(k/\Lambda)}+\bar{c}^a(-k)c^a(k)\frac{k^2}{K(k/\Lambda)}\right]$$

Both the Ward identities and ghost equations of motion come from the invariance of the action $\mathcal{S}$:

1. **BRST invariance**

$$\begin{aligned}\Sigma\ \equiv\ &\int_k K(k/\Lambda)\Bigg[\frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(-k)}\mathcal{S}\cdot\frac{\delta\mathcal{S}}{\delta A^a_\mu(k)}+\frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(-k)}\frac{\delta\mathcal{S}}{\delta A^a_\mu(k)}\\&+\frac{\delta\mathcal{S}}{\delta\bar{c}^{a*}(k)}\frac{\overrightarrow{\delta}}{\delta\bar{c}^a(-k)}\mathcal{S}+\frac{\delta}{\delta\bar{c}^{a*}(k)}\frac{\overrightarrow{\delta}}{\delta\bar{c}^a(-k)}\mathcal{S}\\&-\frac{\delta\mathcal{S}}{\delta c^{a*}(-k)}\cdot\mathcal{S}\frac{\overleftarrow{\delta}}{\delta c^a(k)}-\frac{\delta\mathcal{S}}{\delta c^{a*}(-k)}\frac{\overleftarrow{\delta}}{\delta c^a(k)}\Bigg]=0\end{aligned}$$

This is the “quantum” invariance of the action under the BRST transformation:

$$\begin{cases}\delta_\epsilon A^a_\mu(k)&=&K(k/\Lambda)\epsilon\frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(-k)}\mathcal{S}\\ \delta_\epsilon\bar{c}^a(-k)&=&K(k/\Lambda)\epsilon\frac{\delta\mathcal{S}}{\delta\bar{c}^{a*}(k)}\\ \delta_\epsilon c^a(k)&=&K(k/\Lambda)\epsilon\frac{\delta\mathcal{S}}{\delta c^{a*}(-k)}\end{cases}$$

At the vanishing sources, we obtain

$$\left\{ \begin{array}{rcl} \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \mathcal{S} & = & \frac{1}{i} \left[(D_\mu c)^a \right] = k_\mu \mathcal{C}^a(k) + \left[A_\mu c \right]^a (k) \\ \frac{\delta \mathcal{S}}{\delta \bar{c}^{a*}(k)} & = & -\frac{1}{\xi} k_\mu \mathcal{A}_\mu^a(-k) \\ \frac{\delta \mathcal{S}}{\delta c^{a*}(-k)} & = & \left[cc \right]^a (k) \end{array} \right.$$

where $[A_\mu c]^a$ is the composite operator corresponding to

$$igf^{abc} A_\mu^b c^c$$

and $[cc]$ corresponds to

$$\frac{g}{2i} f^{abc} c^b c^c$$

$\mathcal{C}^a$ and $\mathcal{A}_\mu^a$ are the composite operators corresponding to the elementary fields c^a, A_μ^a:

$$\left\{ \begin{array}{rcl} \mathcal{C}^a(k) & \equiv & c^a(k) + \frac{1-K(k/\Lambda)}{k^2} \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} \mathcal{S}_{\mathrm{int}} \\ \mathcal{A}_\mu^a(k) & \equiv & A_\mu^a(k) + \frac{1-K(k/\Lambda)}{k^2} \left(\delta_{\mu\nu} - (1-\xi) \frac{k_\mu k_\nu}{k^2} \right) \frac{\delta \mathcal{S}_{\mathrm{int}}}{\delta A_\nu^a(-k)} \end{array} \right.$$

2. **The ghost equation of motion**

$$k_\mu \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \mathcal{S} = -K(k/\Lambda) \frac{\overrightarrow{\delta}}{\delta \bar{c}^a(-k)} \mathcal{S}$$

This is an equality between two composite operators.

The ghost equation implies that the dependence of $\mathcal{S}$ on A_μ^{a*} and $\bar{c}^a$ is only through the linear combination

$$A_\mu^{a*}(-k) - \frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k)$$

Using this we can simplify the BRST invariance as follows:

$$\begin{aligned} \Sigma \quad \equiv \quad & \int_k K(k/\Lambda) \Bigg[\frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \mathcal{S} \cdot \left(\frac{\delta \mathcal{S}}{\delta A_\mu^a(k)} - \frac{1}{K(k/\Lambda)} k_\mu \frac{\delta \mathcal{S}}{\delta \bar{c}^{a*}(k)} \right) \\ & + \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \left(\frac{\delta \mathcal{S}}{\delta A_\mu^a(k)} - \frac{1}{K(k/\Lambda)} k_\mu \frac{\delta \mathcal{S}}{\delta \bar{c}^{a*}(k)} \right) \\ & - \frac{\delta \mathcal{S}}{\delta c^{a*}(-k)} \cdot \mathcal{S} \frac{\overleftarrow{\delta}}{\delta c^a(k)} - \frac{\delta \mathcal{S}}{\delta c^{a*}(-k)} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \Bigg] = 0 \end{aligned}$$

As in QED, the source $-\frac{1}{\xi} k_\mu \bar{c}^{a*}(k)$ couples to the elementary field $A_\mu^a(-k)$. Hence, the dependence of the action on $\bar{c}^{a*}$ is obtained as a shift of A_μ^a, and the action can be written as

$$\begin{aligned}
\mathcal{S} \quad = \quad & -\frac{1}{2}\int_k A^a_\mu(k) A^a_\nu(-k) \frac{k^2\delta_{\mu\nu} - \left(1-\frac{1}{\xi}\right) k_\mu k_\nu}{K(k/\Lambda)} \\
& + \int_k \left(A^{a*}_\mu(-k) - \frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k) \right) k_\mu c^a(k) \\
& -\frac{1}{\xi}\int_k k_\mu \bar{c}^{a*}(k) A^a_\mu(-k) - \frac{1}{2\xi}\int_k \left(1-K(k/\Lambda)\right) \bar{c}^{a*}(-k)\bar{c}^{a*}(k) \\
& +\mathcal{S}_{\rm int}\Bigg[A^a_\mu(k) - \frac{1-K(k/\Lambda)}{k^2} k_\mu \bar{c}^{a*}(k), A^{a*}_\mu(-k) - \frac{1}{K(k/\Lambda)} k_\mu \bar{c}^a(-k), \\
& \qquad\qquad c^a(k), c^{a*}(-k) \Bigg]
\end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
k_\mu \frac{\delta \mathcal{S}}{\delta \bar{c}^{a*}(k)}\Big|_{\bar{c}*=0} \quad &= \quad -\frac{1}{\xi} k_\mu k_\nu A^a_\nu(-k) - \frac{1-K(k/\Lambda)}{k^2} k_\mu k_\nu \frac{\delta \mathcal{S}_{\rm int}}{\delta A^a_\nu(k)}\Big|_{\bar{c}*=0} \\
k_\mu \frac{\delta \mathcal{S}}{\delta A^a_\mu(k)}\Big|_{\bar{c}^*=0} \quad &= \quad -\frac{1}{\xi}\frac{1}{K(k/\Lambda)} k^2 k_\nu A^a_\nu(-k) + k_\mu \frac{\delta \mathcal{S}_{\rm int}}{\delta A^a_\mu(k)}\Big|_{\bar{c}^*=0}
\end{aligned}$$

Hence, we find

$$k_\mu \frac{\delta \mathcal{S}}{\delta \bar{c}^{a*}(k)}\Big|_{\bar{c}*=0} = -\frac{1-K(k/\Lambda)}{k^2} k_\mu k_\nu \frac{\delta \mathcal{S}}{\delta A^a_\nu(k)}\Big|_{\bar{c}^*=0} - \frac{1}{\xi}\frac{1}{K(k/\Lambda)} k_\mu k_\nu A^a_\nu(-k)$$

Thus, we obtain

$$\begin{aligned}
& \frac{\delta \mathcal{S}}{\delta A^a_\mu(k)}\Big|_{\bar{c}*=0} - \frac{1}{K(k/\Lambda)} k_\mu \frac{\delta \mathcal{S}}{\delta \bar{c}^{a*}(k)}\Big|_{\bar{c}*=0} \\
& = \left(\delta_{\mu\nu} + \frac{1-K(k/\Lambda)}{K(k/\Lambda)} \frac{k_\mu k_\nu}{k^2} \right) \frac{\delta \mathcal{S}}{\delta A^a_\nu(k)}\Big|_{\bar{c}*=0} + \frac{1}{\xi}\frac{1}{K(k/\Lambda)^2} k_\mu k_\nu A^a_\nu(-k) \\
& = \left(\delta_{\mu\nu} + \frac{1-K(k/\Lambda)}{K(k/\Lambda)} \frac{k_\mu k_\nu}{k^2} \right) \frac{\delta}{\delta A^a_\nu(k)} \left(\mathcal{S} + \frac{1}{2\xi}\int_k \frac{k_\mu k_\nu}{K(k/\Lambda)} A^a_\mu(k) A^a_\nu(-k) \right)\Big|_{\bar{c}^*=0}
\end{aligned}$$

Therefore, taking $\bar{c}=0$ and $\bar{c}^*=0$, the BRST invariance gives

$$\begin{aligned}
\Sigma \equiv \int_k K(k/\Lambda) \Bigg[& \frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(-k)} \mathcal{S}' \cdot P_{\mu\nu}(k) \frac{\delta \mathcal{S}'}{\delta A^a_\nu(k)} + \frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(-k)} P_{\mu\nu}(k) \frac{\delta \mathcal{S}'}{\delta A^a_\nu(k)} \\
& - \frac{\delta \mathcal{S}'}{\delta c^{a*}(-k)} \cdot \mathcal{S}' \frac{\overleftarrow{\delta}}{\delta c^a(k)} - \frac{\delta \mathcal{S}'}{\delta c^{a*}(-k)} \frac{\overleftarrow{\delta}}{\delta c^a(k)} \Bigg] = 0
\end{aligned}$$

where we have defined

$$P_{\mu\nu}(k) \equiv \delta_{\mu\nu} + \frac{1-K(k/\Lambda)}{K(k/\Lambda)} \frac{k_\mu k_\nu}{k^2}$$

and

$$\begin{aligned}
\mathcal{S}' &\equiv \mathcal{S}_{\bar{c}^*=0} + \frac{1}{2\xi}\int_k \frac{k_\mu k_\nu}{K(k/\Lambda)} A^a_\mu(k) A^a_\nu(-k) \\
&= -\frac{1}{2}\int_k A^a_\mu(k)A^a_\nu(-k)\frac{k^2\delta_{\mu\nu}-k_\mu k_\nu}{K(k/\Lambda)} + \int_k A^{a*}_\mu(-k)k_\mu c^a(k) \\
&\quad + \mathcal{S}_{\mathrm{int}}\left[A^a_\mu(k), A^{a*}_\mu(-k), c^a(k), c^{a*}(-k)\right]
\end{aligned}$$

Once we obtain $\mathcal{S}'$, we can construct $\mathcal{S}$ for the non-vanishing $\bar{c}, \bar{c}^*$, since we know how the action $\mathcal{S}$ depends on $\bar{c}$ and $\bar{c}^*$. We will write $\mathcal{S}'$ as $\mathcal{S}$ from now on.

Our task is now to obtain $\mathcal{S}$ that satisfies the BRST invariance given above. From now on, we only have A^a_μ, A^{a*}_μ and c^a, c^{a*} as fields and sources. Let us define δ_Q by

$$\begin{aligned}
\delta_Q \equiv &\int_k K(k/\Lambda)\Bigg[P_{\mu\nu}(k)\frac{\delta\mathcal{S}}{\delta A^a_\nu(k)}\cdot\frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(k)} \\
&+\frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(k)}\mathcal{S}\cdot P_{\mu\nu}(k)\frac{\delta}{\delta A^a_\nu(k)} + \frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(k)}P_{\mu\nu}(k)\frac{\delta}{\delta A^a_\nu(k)} \\
&-\mathcal{S}\frac{\overleftarrow{\delta}}{\delta c^a(k)}\cdot\frac{\delta}{\delta c^{a*}(-k)} + \frac{\delta\mathcal{S}}{\delta c^{a*}(-k)}\cdot\frac{\overrightarrow{\delta}}{\delta c^a(k)} + \frac{\delta}{\delta c^{a*}(-k)}\frac{\overrightarrow{\delta}}{\delta c^a(k)}\Bigg] = 0
\end{aligned}$$

This is nilpotent under the assumption $\Sigma = 0$.
HW#17: Show this is the case. (Hint: In general δ_Q defined by

$$\delta_Q\mathcal{O} \equiv \mathrm{e}^{-\mathcal{S}}\int_q f_{ij}(q)\sum_i \frac{\overrightarrow{\delta}}{\delta\phi_i(q)}\frac{\overrightarrow{\delta}}{\delta\phi^*_j(-q)}\left(\mathrm{e}^{\mathcal{S}}\mathcal{O}\right)$$

is nilpotent if it is fermionic.)

By the definition of δ_Q and Σ, we obtain the algebraic constraint:

$$\begin{aligned}
\delta_Q\Sigma \equiv &\int_k K(k/\Lambda)\Bigg[P_{\mu\nu}(k)\frac{\delta\mathcal{S}}{\delta A^a_\nu(k)}\cdot\frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(k)}\Sigma \\
&+\frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(k)}\mathcal{S}\cdot P_{\mu\nu}(k)\frac{\delta}{\delta A^a_\nu(k)}\Sigma + \frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu(k)}P_{\mu\nu}(k)\Sigma\frac{\delta}{\delta A^a_\nu(k)} \\
&-\mathcal{S}\frac{\overleftarrow{\delta}}{\delta c^a(k)}\cdot\frac{\delta\Sigma}{\delta c^{a*}(-k)} - \frac{\delta\mathcal{S}}{\delta c^{a*}(-k)}\cdot\Sigma\frac{\overleftarrow{\delta}}{\delta c^a(k)} - \frac{\delta}{\delta c^{a*}(-k)}\Sigma\frac{\overleftarrow{\delta}}{\delta c^a(k)}\Bigg] = 0
\end{aligned}$$

Now, our task is to construct the loop expansion

$$\mathcal{S} = \sum_{l=0}^{\infty}\mathcal{S}_l$$

Assuming that the BRST invariance holds up to $(l-1)$-loop:

$$\Sigma_0 = \cdots = \Sigma_{l-1} = 0$$

we wish to construct $\mathcal{S}_l$ by fine tuning the parameters of the theory so that

$$\Sigma_l = 0$$

is satisfied. The question is if we have enough parameters to satisfy this condition. To answer this, we need the help of the algebraic constraint $(\delta_Q \Sigma)_l = 0$.

At l-loop, the algebraic constraint gives

$$\begin{aligned} (\delta_Q \Sigma)_l \quad \equiv \quad & \int_k K(k/\Lambda) \Bigg[P_{\mu\nu}(k) \frac{\delta \mathcal{S}_0}{\delta A_\nu^a(k)} \cdot \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \Sigma_l \\ & + \frac{\overrightarrow{\delta}}{\delta A_\mu^{a*}(-k)} \mathcal{S}_0 \cdot P_{\mu\nu}(k) \frac{\delta \Sigma_l}{\delta A_\nu^a(k)} \\ & - \mathcal{S}_0 \frac{\overleftarrow{\delta}}{\delta c^a(k)} \cdot \frac{\delta \Sigma_l}{\delta c^{a*}(-k)} - \frac{\delta \mathcal{S}_0}{\delta c^{a*}(-k)} \cdot \Sigma_l \frac{\overleftarrow{\delta}}{\delta c^a(k)} \Bigg] = 0 \end{aligned}$$

We note that the second order differentials in the integrand do not contribute since $\Sigma_{l-1} = 0$. (If $\Sigma_{l-1} \neq 0$, it would have contributed to $(\delta_Q \Sigma)_l$, since the integral over q gives an extra loop.) Clearly, the algebraic constraint restricts the possible form of Σ_l.

As a preparation, let us tabulate the basic properties of the fields[15]:

field	statistics	dimension	ghost number
A_μ^a	b	1	0
c^a	f	1	1
A_μ^{a*}	f	2	-1
c^{a*}	b	2	-2

From now on, we examine the case of SU(2) for which the structure constant is the totally antisymmetric tensor:

$$f^{abc} = \epsilon^{abc}$$

Now, let us look at the asymptotic behavior of $\mathcal{S}_l$ at a large cutoff. Using the coordinate space notation for convenience, the asymptotic behavior of $\mathcal{S}_l$ is given by

$$\begin{aligned} \mathcal{S}_l \quad \to \quad & \int d^4x \Bigg[a_1 \frac{1}{2} (\partial_\mu A_\nu^a)^2 + a_2 \frac{1}{2} (\partial_\mu A_\mu^a)^2 + a_3 g \epsilon^{abc} \partial_\mu A_\nu^a A_\mu^b A_\nu^c \\ & + a_4 \frac{g^2}{4} A_\mu^a A_\mu^a A_\nu^b A_\nu^b + a_5 \frac{g^2}{4} A_\mu^a A_\nu^a A_\mu^b A_\nu^b \\ & + a_6 A_\mu^{a*} \partial_\mu c^a + a_7 g \epsilon^{abc} A_\mu^{a*} A_\mu^b c^c + a_8 \frac{g}{2} \epsilon^{abc} c^{a*} c^b c^c \Bigg] \end{aligned}$$

---

[15] We do not need to consider $\bar{c}^a$, since the dependence on $\bar{c}^a$ is determined by that on A_μ^*.

where we have ignored the terms proportional to Λ^2 since they are completely determined by the ERG differential equation. The eight coefficients $a_1, \cdots, a_8$ are functions of $\ln \Lambda/\mu$ whose values at $\Lambda = \mu$ are free to choose. The above $\mathcal{S}_l$ is the most general bosonic expression with zero ghost number. Especially, for $l = 0$, we choose the following asymptotic form

$$\begin{aligned}\mathcal{S}_0 \quad \to \quad & \int d^4x \left[\frac{1}{2}(\partial_\mu A^a_\nu)^2 - \frac{1}{2}(\partial_\mu A^a_\mu)^2 \right. \\ & + g\epsilon^{abc}\partial_\mu A^a_\nu A^b_\mu A^c_\nu + \frac{g^2}{4}\left(A^a_\mu A^a_\mu A^b_\nu A^b_\nu - A^a_\mu A^a_\nu A^b_\mu A^b_\nu \right) \\ & \left. + A^{a*}_\mu \frac{1}{i}\left(\partial_\mu c^a - g\epsilon^{abc} A^b_\mu c^c\right) + c^{a*}\frac{g}{2i}\epsilon^{abc} c^b c^c \right]\end{aligned}$$

as the starting point of the perturbative construction.

We next consider the asymptotic behavior of Σ_l. This is a dimension 5 operator with ghost number 1. There are eleven terms:

$$\begin{aligned}\Sigma_l \quad \to \quad & \int d^4x \left[s_1 c^a \partial^2 \partial_\mu A^a_\mu + s_2 g\epsilon^{abc} c^a \partial^2 A^b_\mu \cdot A^c_\mu + s_3 g\epsilon^{abc} c^a \partial_\mu \partial_\nu A^b_\nu \cdot A^c_\mu \right. \\ & + s_4 g^2 c^a \partial_\mu A^a_\mu \cdot A^b_\nu A^b_\nu + s_5 g^2 c^a \partial_\mu A^a_\nu \cdot A^b_\mu A^b_\nu + s_6 g^2 c^a A^a_\mu \partial_\nu A^b_\nu A^b_\mu \\ & + s_7 g^2 c^a A^a_\mu \partial_\nu A^b_\mu A^b_\nu + s_8 g^2 c^a A^a_\mu \partial_\mu A^b_\nu A^b_\nu \\ & \left. + s_9 g\epsilon^{abc} c^a c^b \partial_\mu A^{c*}_\mu + s_{10} g\epsilon^{abc} \partial_\mu c^a c^b A^{c*}_\mu + s_{11} g^2 c^a c^b A^{a*}_\mu A^a_\mu \right]\end{aligned}$$

At first sight we may not be able to make this vanish using only eight parameters. But the eleven parameters are not all independent, since $(\delta_Q \Sigma)_l$ must vanish. Looking at the asymptotic behavior of $(\delta_Q \Sigma)_l$, we obtain

$$\begin{aligned}(\delta_Q \Sigma)_l \quad \to \quad & \int d^4x \left[\left(\frac{\delta \mathcal{S}_0}{\delta A^a_\mu} - i\partial_\mu \frac{\delta \mathcal{S}_0}{\delta \bar{c}^{a*}} \right) \frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu} \Sigma_l + \frac{\overrightarrow{\delta}}{\delta A^{a*}_\mu} \mathcal{S}_0 \cdot \left(\frac{\delta}{\delta A^a_\mu} - i\partial_\mu \frac{\delta}{\delta \bar{c}^{a*}} \right) \Sigma_l \right. \\ & \left. - \mathcal{S}_0 \frac{\overleftarrow{\delta}}{\delta c^a} \frac{\delta \Sigma_l}{\delta c^{a*}} - \frac{\delta \mathcal{S}_0}{\delta c^{a*}} \cdot \Sigma_l \frac{\overleftarrow{\delta}}{\delta c^a} \right] = 0\end{aligned}$$

This constrains the s parameters. Becchi has shown the possibility of satisfying the BRST invariance in his Parma lectures.[3] I hope to provide more details in an updated version of the present notes. ;-)

Appendix: Introduction of auxiliary fields B^a

Instead of using the gauge fixing term

$$-\frac{1}{2\xi}\int_k A_\mu^a(k)A_\nu^a(-k)\frac{1}{K(k/\Lambda)}k_\mu k_\nu$$

we can gauge fix the theory using auxiliary fields B^a coupled to $\frac{1}{\xi}k_\mu A_\mu^a$. B^a do not transform under the BRST, and we do not introduce their antifields. The full action can now be written in the following form:

$$\begin{aligned}
\mathcal{S} \quad = \quad & -\frac{1}{2}\int_k A_\mu^a(k)A_\nu^a(-k)\frac{1}{K(k/\Lambda)}\left(k^2\delta_{\mu\nu}-k_\mu k_\nu\right) \\
& +\int_k \frac{1}{K(k/\Lambda)}B^a(-k)k_\mu A_\mu^a(k) - \frac{\xi}{2}\int_k \frac{1}{K(k/\Lambda)}B^a(-k)B^a(k) \\
& +\int_k \left(A_\mu^{a*}(-k) - \frac{1}{K(k/\Lambda)}k_\mu \bar{c}^a(-k)\right)k_\mu c^a(k) + \int_k \bar{c}^{a*}(k)B^a(-k) \\
& +\mathcal{S}_{\mathrm{int}}\Bigg[A_\mu^a(k) - \frac{1-K(k/\Lambda)}{k^2}k_\mu \bar{c}^{a*}(k), A_\mu^{a*}(-k) - \frac{1}{K(k/\Lambda)}k_\mu \bar{c}^a(-k), \\
& \qquad c^a(k), c^{a*}(-k)\Bigg]
\end{aligned}$$

Substituting this into the BRST invariance $\Sigma = 0$ (Σ is defined the same way as before since there is no antifield for B), and setting $\bar{c} = \bar{c}^* = B = 0$, we find that $\mathcal{S}$ satisfies the same BRST invariance as $\mathcal{S}'$ that we have obtained without using B. Thus, there is nothing to gain or lose by introducing the auxiliary fields.

5 Non-perturbative aspects of the Wilson ERG

So far our discussions of ERG have been restricted to perturbation theory. Actually, Ken Wilson introduced ERG[16] to define the continuum limits in quantum field theory non-perturbatively. In this final part of the lectures we wish to discuss the non-perturbative aspects of the ERG.[17] We will first introduce a general framework for ERG differential equations to relate Wilson's ERG differential equation to Polchinski's. We then apply ERG to define the non-trivial continuum limit of the scalar theory in $D = 3$.

5.1 Generalized diffusion equation

We consider a real scalar theory in D dimensional euclidean space. Given an action $S[\phi]$, we construct a new action $S_t[\phi]$ by the following integral transformation:

$$\begin{aligned}
&\exp\left[S_t[\phi]\right]\\
&\equiv \int [d\phi'] \exp\left[-\frac{1}{2}\int_p \left(A_t(p)\phi(p) - B_t(p)\phi'(p)\right)\left(A_t(p)\phi(-p) - B_t(p)\phi'(-p)\right) + S[\phi']\right]\\
&= \int [d\phi'] \exp\left[-\frac{1}{2}\int_p A_t(p)^2 \left(\phi(p) - \frac{B_t(p)}{A_t(p)}\phi'(p)\right)\left(\phi(-p) - \frac{B_t(p)}{A_t(p)}\phi'(-p)\right) + S[\phi']\right]
\end{aligned}$$

where we assume both A_t and B_t are functions of p^2. The physical meaning of this transformation is clear: $\phi(p)$ is diffused around

$$\frac{B_t(p)}{A_t(p)}\phi'(p)$$

with the width $\frac{1}{A_t(p)}$. The distribution of ϕ' is given in terms of the weight $\mathrm{e}^{S[\phi']}$. This is analogous to the solution of the diffusion equation

$$\partial_t P(t,x) = D\partial_x^2 P(t,x)$$

for the distribution function $P(t,x)$ in the coordinate space at time t. The solution is given by the integral formula

$$P(t,x) = \sqrt{\frac{4Dt}{\pi}} \int_{-\infty}^{\infty} dy\, \exp\left[-\frac{1}{4Dt}(x-y)^2\right] P(0,y)$$

We now wish to derive the differential equation for $S_t[\phi]$ analogous to the above diffusion equation. For this, we change variables to rewrite

$$\begin{aligned}
&\exp\left[S_t[\phi]\right]\\
&= \int [d\phi'] \exp\left[-\frac{1}{2}\int_p \phi'(p)\phi'(-p) + S\left[\frac{1}{B_t(p)}\phi'(p) + \frac{A_t(p)}{B_t(p)}\phi(p)\right]\right]
\end{aligned}$$

---

[16] He called it simply RG.

[17] We will review the materials explained in sects. 11 and 12 of the Wilson-Kogut lecture notes.[1]

Differentiating this with respect to t, we obtain

$$\begin{aligned}
&\partial_t S_t[\phi]\,\exp[S_t[\phi]]\\
&= \int_p \int [d\phi'] \exp\left[-\frac{1}{2}\int_p \phi'(p)\phi'(-p)\right] \left(\partial_t \frac{1}{B_t(p)}\cdot\phi'(p) + \partial_t \frac{A_t(p)}{B_t(p)}\phi(p)\right)\\
&\qquad\qquad \times \frac{B_t(p)}{A_t(p)} \frac{\delta}{\delta\phi(p)} \exp\left[S\left[\frac{1}{B_t(p)}\phi'(p) + \frac{A_t(p)}{B_t(p)}\phi(p)\right]\right]\\
&= \int_p \frac{B_t(p)}{A_t(p)} \partial_t \frac{1}{B_t(p)} \cdot \frac{\delta}{\delta\phi(p)} \int [d\phi']\phi'(p) \exp\left[-\frac{1}{2}\int_p \phi'(p)\phi'(-p) + S\right]\\
&\quad + \int_p \partial_t \ln\frac{A_t(p)}{B_t(p)} \cdot \phi(p) \frac{\delta S_t[\phi]}{\delta\phi(p)} \exp\left[S_t[\phi]\right]
\end{aligned}$$

Since

$$\int [d\phi'] \frac{\delta}{\delta\phi'(-p)} \exp\left[-\frac{1}{2}\int_p \phi'(p)\phi'(-p) + S\left[\frac{1}{B_t(p)}\phi'(p) + \frac{A_t(p)}{B_t(p)}\phi(p)\right]\right] = 0$$

we obtain

$$\int [d\phi']\phi'(p) \exp\left[-\frac{1}{2}\int_p \phi'\phi' + S\right] = \frac{1}{A_t(p)} \frac{\delta S_t[\phi]}{\delta\phi(-p)} \exp\left[S_t[\phi]\right]$$

Hence, we obtain

$$\begin{aligned}
\partial_t S_t[\phi] \quad = \quad & \int_p \left[\frac{B_t(p)}{A_t(p)^2}\partial_t \frac{1}{B_t(p)} \cdot \left\{\frac{\delta S_t[\phi]}{\delta\phi(p)}\frac{\delta S_t[\phi]}{\delta\phi(-p)} + \frac{\delta^2 S_t[\phi]}{\delta\phi(p)\delta\phi(-p)}\right\}\right.\\
&\qquad \left. + \partial_t \ln\frac{A_t(p)}{B_t(p)} \cdot \phi(p)\frac{\delta S_t[\phi]}{\delta\phi(p)}\right]
\end{aligned}$$

Defining

$$\left\{\begin{array}{rcl}
F_t(p) & \equiv & \partial_t \ln \frac{A_t(p)}{B_t(p)}\\
G_t(p) & \equiv & 2\frac{B_t(p)}{A_t(p)^2}\partial_t \frac{1}{B_t(p)} = -2\left\{-\partial_t \ln\frac{A_t(p)}{B_t(p)} + \frac{\partial_t A_t(p)}{A_t(p)}\right\}\frac{1}{A_t(p)^2}
\end{array}\right.$$

we can write

$$\begin{aligned}
\partial_t S_t \quad = \quad & \int_p \left[F_t(p)\cdot\phi(p)\frac{\delta S_t}{\delta\phi(p)}\right.\\
&\qquad \left. + G_t(p)\cdot\frac{1}{2}\left\{\frac{\delta S_t}{\delta\phi(p)}\frac{\delta S_t}{\delta\phi(-p)} + \frac{\delta^2 S_t}{\delta\phi(p)\delta\phi(-p)}\right\}\right]
\end{aligned}$$

This has the familiar form of the ERG differential equation.

We now define the generating functionals of connected correlation functions:

$$\left\{\begin{array}{rcl}
\mathrm{e}^{W[J]} & = & \int [d\phi] \exp\left[S[\phi] + i\int_p J(p)\phi(-p)\right]\\
\mathrm{e}^{W_t[J]} & = & \int [d\phi] \exp\left[S_t[\phi] + i\int_p J(p)\phi(-p)\right]
\end{array}\right.$$

We can compute $W_t[J]$ in terms of $W[J]$ as follows:

$$\begin{aligned}
\mathrm{e}^{W_t[J]} &= \int [d\phi] \mathrm{e}^{i\int_p J(p)\phi(-p)} \int [d\phi'] \exp\Bigg[\\
&\quad -\frac{1}{2}\int_p \left(A_t(p)\phi(p) - B_t(p)\phi'(p)\right)\left(A_t(p)\phi(-p) - B_t(p)\phi'(-p)\right) + S[\phi']\Bigg] \\
&= \int [d\phi] \int [d\phi'] \exp\Bigg[i\int_p \frac{J(p)}{A_t(p)} \left(A_t(p)\phi(-p) - B_t(p)\phi'(-p)\right) \\
&\quad -\frac{1}{2}\int_p \left(A_t(p)\phi(p) - B_t(p)\phi'(p)\right)\left(A_t(p)\phi(-p) - B_t(p)\phi'(-p)\right) \\
&\quad + S[\phi'] + i\int_p J(p)\frac{B_t(p)}{A_t(p)}\phi'(-p)\Bigg] \\
&= \int [d\phi] \exp\left[-\frac{1}{2}\int_p \phi(p)\phi(-p) + i\int_p \frac{J(p)}{A_t(p)}\phi(-p)\right] \\
&\quad \times \int [d\phi'] \exp\left[S[\phi'] + i\int_p J(p)\frac{B_t(p)}{A_t(p)}\phi'(-p)\right] \\
&= \exp\left[-\frac{1}{2}\int_p \frac{1}{A_t(p)^2} J(p)J(-p) + W\left[\frac{B_t(p)}{A_t(p)}J(p)\right]\right]
\end{aligned}$$

This implies

$$\left\{ \begin{array}{rcl} \langle\phi(p)\phi(-p)\rangle_{S_t} &=& \frac{1}{A_t(p)^2} + \frac{B_t(p)^2}{A_t(p)^2}\langle\phi(p)\phi(-p)\rangle_S \\ \langle\phi(p_1)\cdots\phi(p_n)\rangle_{S_t}^{\text{connected}} &=& \prod_{i=1}^n \frac{B_t(p_i)}{A_t(p_i)} \cdot \langle\phi(p_1)\cdots\phi(p_n)\rangle_S^{conn.} \quad (n>1) \end{array} \right.$$

In all the examples we will discuss below, we choose

$$\boxed{F_t(p) \equiv \partial_t \ln \frac{A_t(p)}{B_t(p)} = \frac{\Delta(p\mathrm{e}^t)}{K(p\mathrm{e}^t)} - \frac{\eta(t)}{2}}$$

where $\Lambda = \mu \mathrm{e}^{-t}$ is the cutoff scale, and we have chosen $\mu = 1$ for convenience. Hence,

$$\boxed{\frac{A_t(p)}{B_t(p)} = \frac{K(p)}{K(p\mathrm{e}^t)} \exp\left[-\frac{1}{2}\int_0^t dt'\,\eta(t')\right]}$$

Note we have chosen

$$\frac{A_0(p)}{B_0(p)} = 1$$

For

$$S_0 = S$$

we must also choose

$$\frac{1}{A_0(p)} = 0$$

With our choice for $F_t(p)$, we obtain

$$\begin{array}{rcl} G_t(p) & = & -2\left\{-F_t(p) + \partial_t \ln A_t(p)\right\} \dfrac{1}{A_t(p)^2} \\ & = & -2\left\{-\dfrac{\Delta(p\mathrm{e}^t)}{K(p\mathrm{e}^t)} + \dfrac{\eta}{2} + \partial_t \ln A_t(p)\right\} \dfrac{1}{A_t(p)^2} \\ & = & -2\partial_t \ln\left(A_t(p)\mathrm{e}^{\int^t \frac{\eta}{2}} \dfrac{K(p\mathrm{e}^t)}{K(p)}\right) \dfrac{1}{A_t(p)^2} \end{array}$$

Hence, defining

$$\bar{A}_t(p) \equiv A_t(p)\mathrm{e}^{\int^t \frac{\eta}{2}} \frac{K(p\mathrm{e}^t)}{K(p)}$$

we obtain

$$G_t(p) = -2\mathrm{e}^{\int^t \eta} \frac{K(p\mathrm{e}^t)^2}{K(p)^2} \frac{\partial_t \ln \bar{A}_t(p)}{\bar{A}_t(p)^2}$$

Hence,

$$\partial_t \frac{1}{\bar{A}_t(p)^2} = \mathrm{e}^{-\int^t \eta} \frac{K(p)^2}{K(p\mathrm{e}^t)^2} G_t(p)$$

Let us now look at two examples of $G_t(p)$ and calculate the corresponding $A_t(p)$.

Example 1 (Wilson)

We choose

$$G_t(p) = 2\mathrm{e}^{2t}\left(\frac{\Delta(p\mathrm{e}^t)}{K(p\mathrm{e}^t)} + 1 - \frac{1}{2}\eta(t)\right)$$

This gives

$$\begin{array}{rcl} \partial_t \dfrac{1}{\bar{A}_t(p)^2} & = & 2\mathrm{e}^{2t-\int^t \eta} \dfrac{K(p)^2}{K(p\mathrm{e}^t)^2} \left(\dfrac{\Delta(p\mathrm{e}^t)}{K(p\mathrm{e}^t)} + 1 - \dfrac{\eta}{2}\right) \\ & = & \partial_t \left(\mathrm{e}^{2t-\int^t \eta} \dfrac{K(p)^2}{K(p\mathrm{e}^t)^2}\right) \end{array}$$

Hence, we obtain

$$\frac{1}{\bar{A}_t(p)^2} = \mathrm{e}^{2t-\int^t \eta} \frac{K(p)^2}{K(p\mathrm{e}^t)^2} - 1$$

using the initial condition $1/A_0 = 0$. Thus, we obtain

$$\frac{1}{A_t(p)^2} = \mathrm{e}^{2t} - \frac{K(p\mathrm{e}^t)^2}{K(p)^2}\,\mathrm{e}^{\int_0^t \eta}$$

This implies

$$\left\{\begin{array}{rcl} \langle\phi(p)\phi(-p)\rangle_{S_t} - \mathrm{e}^{2t} & = & \mathrm{e}^{\int_0^t \eta}\, \frac{K(p\mathrm{e}^t)^2}{K(p)^2} \left(\langle\phi(p)\phi(-p)\rangle_S - 1\right) \\ \langle\phi(p_1)\cdots\phi(p_{2n})\rangle_{S_t} & = & \mathrm{e}^{n\int_0^t \eta} \prod_{i=1}^{2n} \frac{K(p_i\mathrm{e}^t)}{K(p_i)} \cdot \langle\phi(p_1)\cdots\phi(p_{2n})\rangle_S \end{array}\right.$$

This is the original choice of Wilson discussed in sect. 11 of [1]. In this choice, the two-point function behaves as

$$\langle \phi(p)\phi(-p)\rangle_{S_t} \overset{t\to\infty}{\longrightarrow} \mathrm{e}^{2t}$$

Example 2 (Polchinski)

We choose

$$G_t(p) = \frac{1}{p^2+m^2(t)}\left\{\Delta(p\mathrm{e}^t) - \left(\eta(t) + \frac{b_m(t)}{p^2+m^2(t)}\right) K(p\mathrm{e}^t)\left(1-K(p\mathrm{e}^t)\right)\right\}$$

where

$$b_m(t) \equiv \frac{dm^2(t)}{dt}$$

This gives the Polchinski ERG differential equation if $\eta = 0$ and m^2 is t-independent.

Let us find the corresponding $A_t(p)$. We find

$$\begin{aligned}
\partial_t \frac{1}{\bar{A}_t(p)^2} &= \mathrm{e}^{-\int^t \eta} \frac{K(p)^2}{K(p\mathrm{e}^t)^2} \\
&\quad \times \frac{1}{p^2+m^2(t)}\left\{\Delta(p\mathrm{e}^t) - \left(\eta(t) + \frac{b_m(t)}{p^2+m^2(t)}\right) K(p\mathrm{e}^t)\left(1-K(p\mathrm{e}^t)\right)\right\} \\
&= K(p)^2 \partial_t \left(\frac{\mathrm{e}^{-\int^t \eta}}{p^2+m^2(t)}\left(\frac{1}{K(p\mathrm{e}^t)} - 1\right)\right)
\end{aligned}$$

Hence,

$$\frac{1}{\bar{A}_t(p)^2} = K(p)^2 \left(\frac{\mathrm{e}^{-\int^t \eta}}{p^2+m^2(t)}\left(\frac{1}{K(p\mathrm{e}^t)} - 1\right) - \frac{1}{p^2+m^2(0)}\left(\frac{1}{K(p)} - 1\right)\right)$$

Thus, we obtain

$$\frac{1}{A_t(p)^2} = K(p\mathrm{e}^t)^2 \left\{\frac{1}{p^2+m^2(t)}\left(\frac{1}{K(p\mathrm{e}^t)} - 1\right) - \frac{\mathrm{e}^{\int_0^t \eta}}{p^2+m^2(0)}\left(\frac{1}{K(p)} - 1\right)\right\}$$

Hence, we obtain

$$\begin{aligned}
&\langle \phi(p)\phi(-p)\rangle_{S_t} - \frac{1}{p^2+m^2(t)} K(p\mathrm{e}^t)\left(1-K(p\mathrm{e}^t)\right) \\
&= \mathrm{e}^{\int_0^t \eta} \frac{K(p\mathrm{e}^t)^2}{K(p)^2}\left(\langle \phi(p)\phi(-p)\rangle_S - \frac{1}{p^2+m^2(0)} K(p)(1-K(p))\right)
\end{aligned}$$

and

$$\langle \phi(p_1)\cdots\phi(p_{2n})\rangle_{S_t} = \mathrm{e}^{n\int_0^t \eta} \prod_{i=1}^{2n} \frac{K(p_i\mathrm{e}^t)}{K(p_i)} \cdot \langle \phi(p_1)\cdots\phi(p_{2n})\rangle_S$$

With this choice of $G_t(p)$, we obtain the characteristic asymptotic behavior

$$\langle \phi(p)\phi(-p) \rangle_{S_t} \overset{t\to\infty}{\longrightarrow} \frac{K(p\mathrm{e}^t)}{p^2 + m^2(t)}$$

Before concluding this subsection, let us consider the relation between two different formalisms of ERG: one given by A_t, B_t and another given by C_t, D_t. Let

$$\begin{aligned}
&\exp[S_t[\phi]] \\
&\equiv \int [d\phi'] \exp\left[-\frac{1}{2}(A_t(p)\phi(p) - B_t\phi'(p))(A_t(p)\phi(-p) - B_t(p)\phi'(-p)) + S[\phi'] \right] \\
&\exp[S'_t[\phi]] \\
&\equiv \int [d\phi'] \exp\left[-\frac{1}{2}(C_t(p)\phi(p) - D_t\phi'(p))(C_t(p)\phi(-p) - D_t(p)\phi'(-p)) + S[\phi'] \right]
\end{aligned}$$

We wish to derive the relation between S_t and S'_t. To derive the relation, we recall the following results on the generating functional:

$$\left\{ \begin{array}{rcl} \mathrm{e}^{W_t[J]} &=& \exp\left[W\left[\frac{B_t(p)}{A_t(p)} J(p) \right] - \frac{1}{2}\int_p \frac{1}{A_t(p)^2} J(p)J(-p) \right] \\ \mathrm{e}^{W'_t[J]} &=& \exp\left[W\left[\frac{D_t(p)}{C_t(p)} J(p) \right] - \frac{1}{2}\int_p \frac{1}{C_t(p)^2} J(p)J(-p) \right] \end{array} \right.$$

Hence, we obtain

$$\mathrm{e}^{W'_t[J]} = \exp\left[W_t\left[\frac{A_t D_t}{B_t C_t} J \right] - \frac{1}{2}\int_p J(p)J(-p) \frac{1}{C_t(p)^2}\left(1 - \frac{D_t(p)^2}{B_t(p)^2} \right) \right]$$

This implies

$$\begin{aligned}
&\mathrm{e}^{S'_t[\phi]} = \int [d\phi'] \exp\Bigg[S_t[\phi'] \\
&-\frac{1}{2}\int_p \frac{1}{\frac{1}{C_t(p)^2} - \frac{1}{A_t(p)^2}R_t(p)^2} \left(\phi(p) - R_t(p)\phi'(p)\right)\left(\phi(-p) - R_t(p)\phi'(-p)\right) \Bigg]
\end{aligned}$$

where

$$R_t(p) \equiv \frac{A_t(p)D_t(p)}{B_t(p)C_t(p)}$$

Let us apply this result to the two examples given above. The first example is given by

$$\left\{ \begin{array}{rcl} \frac{1}{A_t(p)^2} &=& \mathrm{e}^{2t} - \frac{K(p\mathrm{e}^t)^2}{K(p)^2}\mathrm{e}^{\int_0^t dt' \eta_W(t')} \\ \frac{A_t(p)}{B_t(p)} &=& \frac{K(p)}{K(p\mathrm{e}^t)}\mathrm{e}^{-\frac{1}{2}\int_0^t dt' \eta_W(t')} \end{array} \right.$$

and the second by

$$\left\{ \begin{array}{rcl} \frac{1}{C_t(p)^2} &=& K(p\mathrm{e}^t)^2 \left\{ \frac{1}{p^2+m^2(t)} \left(\frac{1}{K(p\mathrm{e}^t)} - 1 \right) - \frac{\mathrm{e}^{\int_0^t \eta_P}}{p^2+m^2(0)} \left(\frac{1}{K(p)} - 1 \right) \right\} \\ \frac{C_t(p)}{D_t(p)} &=& \frac{K(p)}{K(p\mathrm{e}^t)}\mathrm{e}^{-\frac{1}{2}\int_0^t dt' \eta_P(t')} \end{array} \right.$$

Thus, we obtain

$$R_t(p) = \mathrm{e}^{\frac{1}{2}\int_0^t dt'\,\left(\eta_P(t')-\eta_W(t')\right)}$$

5.2 Modification of the Polchinski ERG equation

The second example introduced in the previous subsection suggests a possibility of generalizing the Polchinski ERG differential equation:

$$\partial_t S_t = \int_p \frac{\Delta(p\mathrm{e}^t)}{p^2+m^2}\left(\frac{p^2+m^2}{K(p\mathrm{e}^t)}\phi(p)\frac{\delta S_t}{\delta\phi(p)} + \frac{1}{2}\left\{\frac{\delta S_t}{\delta\phi(p)}\frac{\delta S_t}{\delta\phi(-p)} + \frac{\delta^2 S_t}{\delta\phi(p)\delta\phi(-p)}\right\}\right)$$

for which m^2 is a constant, and η vanishes.

Suppose we have a solution $S_t(m^2)$ of the above ERG differential equation. We then modify it infinitesimally by

$$\delta S_t = \frac{\delta z}{2}\mathcal{N} - \delta m^2 \int_p \frac{K(p\mathrm{e}^t)(1-K(p\mathrm{e}^t))}{(p^2+m^2)^2}\frac{1}{2}\left\{\frac{\delta S_t}{\delta\phi(p)}\frac{\delta S_t}{\delta\phi(-p)} + \frac{\delta^2 S_t}{\delta\phi(p)\delta\phi(-p)}\right\}$$

where

$$\mathcal{N} \equiv -\int_p \left(\phi(p)\frac{\delta S_t}{\delta\phi(p)} + 2\frac{K(p\mathrm{e}^t)(1-K(p\mathrm{e}^t))}{p^2+m^2}\frac{1}{2}\left\{\frac{\delta S_t}{\delta\phi(p)}\frac{\delta S_t}{\delta\phi(-p)} + \frac{\delta^2 S_t}{\delta\phi(p)\delta\phi(-p)}\right\}\right)$$

is the composite operator changing the normalization of the field, and the operator proportional to δm^2 is a type 2 operator changing only the two-point function.

The modified action $S_t + \delta S_t$ is no longer a solution of the ERG differential equation, but it is a solution of the ERG differential equation for the modified squared mass $m^2 + \delta m^2$. Moreover, the t-independent correlation functions are the same up to normalization of ϕ.

HW#18: Show that

$$\begin{aligned}
&\frac{1}{K(p\mathrm{e}^t)^2}\left\langle\phi(p)\phi(-p)\right\rangle_{S_t+\delta S_t} + \frac{1-1/K(p\mathrm{e}^t)}{p^2+m^2+\delta m^2}\\
&\quad = (1+\delta z)\left(\frac{1}{K(p\mathrm{e}^t)^2}\left\langle\phi(p)\phi(-p)\right\rangle_{S_t} + \frac{1-1/K(p\mathrm{e}^t)}{p^2+m^2}\right)\\
&\prod_{i=1}^{2n}\frac{1}{K(p_i\mathrm{e}^t)}\cdot\left\langle\phi(p_1)\cdots\phi(p_{2n})\right\rangle_{S_t+\delta S_t}\\
&\quad = (1+n\delta z)\prod_{i=1}^{2n}\frac{1}{K(p_i\mathrm{e}^t)}\cdot\left\langle\phi(p_1)\cdots\phi(p_{2n})\right\rangle_{S_t}
\end{aligned}$$

This proves what is stated in the above paragraph.

We now choose an arbitrary functions $m^2(t')$. Let $S_t(m^2(t'))$ be solutions of the Polchinski ERG differential equation for a fixed squared mass $m^2(t')$. As we



Figure 11: Physically equivalent Polchinski ERG flows with different values of the squared mass parameter. $S_t(m^2(t))$ crosses each trajectory at a single point.

compare $S_t(m^2(t'))$ and $S_t(m^2(t'+\delta t'))$, we can arrange the solutions so that

$$\begin{aligned}\partial_{t'} S_t(m^2(t')) &= \frac{\eta(t')}{2}\mathcal{N}(t')\\ &- \frac{dm^2(t')}{dt'}\int_p \frac{K(pe^t)(1-K(pe^t))}{(p^2+m^2(t'))^2}\frac{1}{2}\left\{\frac{\delta S_t}{\delta\phi(p)}\frac{\delta S_t}{\delta\phi(-p)} + \frac{\delta^2 S_t}{\delta\phi(p)\delta\phi(-p)}\right\}\end{aligned}$$

where $\mathcal{N}(t')$ is the $\mathcal{N}$ operator constructed with $S_t(m^2(t'))$, and $\eta(t')$ is an arbitrary function.

Finally, if we look at the t-dependence of $S_t(m^2(t))$, where t and t' are set equal, we obtain the differential equation of the second example in the previous subsection.

5.3 Renormalization conditions

In this subsection we introduce criteria to choose the t-dependence of the squared mass $m^2(t)$ and the anomalous dimension $\eta(t)$. The criteria, which amount to renormalization conditions, are by no means unique. Let us begin with what is the easiest to understand physically.

1st version

We adopt the following two conditions

$$\begin{cases} \mathcal{V}_2(t;0,0) = 0 \\ \frac{\partial}{\partial p^2}\mathcal{V}_2(t;p,-p)\Big|_{p^2=0} = 0 \end{cases}$$

The second condition is a condition on the normalization of ϕ. Applying the above conditions on the ERG differential equation

$$\begin{aligned}
\partial_t S_t &= \int_p \Bigg[\left(\frac{\Delta(p\mathrm{e}^t)}{K(p\mathrm{e}^t)} - \frac{\eta}{2} \right) \phi(p) \frac{\delta S_t}{\delta\phi(p)} \\
&\quad + \left(\frac{\Delta(p\mathrm{e}^t)}{p^2+m^2(t)} - \left(\eta + \frac{b_m(t)}{p^2+m^2(t)} \right) \frac{K(p\mathrm{e}^t)(1-K(p\mathrm{e}^t))}{p^2+m^2(t)} \right) \\
&\qquad \times \frac{1}{2} \left\{ \frac{\delta S_t}{\delta\phi(p)} \frac{\delta S_t}{\delta\phi(-p)} + \frac{\delta^2 S_t}{\delta\phi(p)\delta\phi(-p)} \right\} \Bigg]
\end{aligned}$$

we obtain

$$\begin{aligned}
-b_m(t) - \eta(t) m^2(t) = \frac{1}{2} \int_q \frac{1}{q^2+m^2(t)} \Bigg\{ & \Delta(q\mathrm{e}^t) \\
& - K(q\mathrm{e}^t)(1-K(q\mathrm{e}^t)) \left(\eta(t) + \frac{b_m(t)}{q^2+m^2(t)} \right) \Bigg\} \mathcal{V}_4(t;q,-q,0,0) \\
-\eta(t) = \frac{1}{2} \frac{\partial}{\partial p^2} \int_q \frac{1}{q^2+m^2(t)} & \left\{ \Delta(q\mathrm{e}^t) - K(q\mathrm{e}^t)(1-K(q\mathrm{e}^t)) \left(\eta(t) + \frac{b_m(t)}{q^2+m^2} \right) \right\} \\
& \times \mathcal{V}_4(t;q,-q,p,-p) \Big|_{p^2=0}
\end{aligned}$$

Thus, we can determine $b_m(t)$ and $\eta(t)$ in terms of the four-point vertex.

2nd version

We introduce a version that is more analogous to the Wilsonian ERG differential equation which has no squared mass parameter. We set to zero the squared mass appearing in the differential equation:

$$m^2(t) = 0$$

Then, the ERG differential equation becomes

$$\begin{aligned}
\partial_t S_t = \int_p \Bigg[& \left(\frac{\Delta(p\mathrm{e}^t)}{K(p\mathrm{e}^t)} - \frac{\eta(t)}{2} \right) \phi(p) \frac{\delta S_t}{\delta\phi(p)} \\
& + \frac{1}{p^2} \left(\Delta(p\mathrm{e}^t) - \eta(t) K(p\mathrm{e}^t)(1-K(p\mathrm{e}^t)) \right) \frac{1}{2} \left\{ \frac{\delta S_t}{\delta\phi(p)} \frac{\delta S_t}{\delta\phi(-p)} + \frac{\delta^2 S_t}{\delta\phi(p)\delta\phi(-p)} \right\} \Bigg]
\end{aligned}$$

We adopt the normalization condition:

$$\frac{\partial}{\partial p^2} \mathcal{V}_2(t;p,-p) \Big|_{p^2=0} = 0$$

This determines $\eta(t)$ as

$$\eta(t) \equiv \frac{-\frac{\partial}{\partial p^2}\int_q \frac{\Delta(q\mathrm{e}^t)}{q^2}\mathcal{V}_4(t;q,-q,p,-p)\Big|_{p^2=0}}{1-\frac{\partial}{\partial p^2}\int_q \frac{K(q\mathrm{e}^t)(1-K(q\mathrm{e}^t))}{q^2}\mathcal{V}_4(t;q,-q,p,-p)\Big|_{p^2=0}}\,,$$

Wilson ERG

The Wilson ERG differential equation is given by

$$\begin{aligned}\partial_t S_t \quad &= \quad \int_p \Bigg[\left\{\left(\frac{\Delta(p\mathrm{e}^t)}{K(p\mathrm{e}^t)}-\frac{\eta(t)}{2}\right)\phi(p)\right\}\frac{\delta S(t)}{\delta\phi(p)} \\ &\qquad + \left(\frac{\Delta(p\mathrm{e}^t)}{K(p\mathrm{e}^t)}+1-\frac{\eta(t)}{2}\right)\left\{\frac{\delta S_t}{\delta\phi(p)}\frac{\delta S_t}{\delta\phi(-p)}+\frac{\delta^2 S_t}{\delta\phi(p)\delta\phi(-p)}\right\}\Bigg]\end{aligned}$$

We denote the functional derivatives of the action as

$$u_{2n}(t;p_1,\cdots,p_{2n})(2\pi)^D\delta^{(D)}(p_1+\cdots+p_{2n}) \equiv \frac{\delta^{2n}S_t}{\delta\phi(p_1)\cdots\phi(p_{2n})}\Big|_{\phi=0}$$

Then, we adopt the normalization condition

$$\frac{\partial}{\partial p^2}u_2(t;p,-p)\Big|_{p^2=0} = -1$$

Applying this to the ERG equation, we obtain

$$\frac{1}{2}\eta(t) = \frac{2u_2(t;0,0)-\frac{1}{2}\frac{\partial}{\partial p^2}\int_q\left(\frac{\Delta(q\mathrm{e}^t)}{K(q\mathrm{e}^t)}+1\right)u_4(t;q,-q,p,-p)\Big|_{p^2=0}}{1+2u_2(t;0,0)-\frac{1}{2}\frac{\partial}{\partial p^2}\int_q u_4(t;q,-q,p,-p)\Big|_{p^2=0}}$$

This depends on both the two- and four-point vertices.

5.4 Rescaling of space

Now, in non-perturbative applications of the ERG, it is the fixed points of the ERG transformation which play the most important roles. A fixed point is an action $S[\phi]$ that does not transform. To define a continuum limit, we need a fixed point. We also need to know the critical exponents characterizing the fixed point.

The ERG transformations we have formulated so far have no fixed point for an obvious reason. The cutoff of the action decreases along the ERG trajectory. For ERG to have fixed points, we must introduce rescaling of space (or momentum) so that the cutoff is fixed under ERG. The purpose of this subsection is to understand how rescaling modifies the ERG differential equations.

The necessary rescaling does not depend on interactions, and it is enough to study the massless free theory:

$$S_t = -\frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2}{K(p\mathrm{e}^t)}$$

which has a cutoff e^{-t}. In order to fix the cutoff at 1, we must measure the momentum p in units of the cutoff. Hence, we must change p to

$$p' \equiv \frac{p}{\mathrm{e}^{-t}} = p\mathrm{e}^t$$

Substituting this into S_t, we obtain

$$S_t = -\frac{1}{2}\int_{p'} \mathrm{e}^{-(D+2)t}\phi(p'\mathrm{e}^{-t})\phi(-p'\mathrm{e}^{-t})\frac{{p'}^2}{K(p')}$$

We now define

$$\phi'(p') \equiv \mathrm{e}^{-\frac{D+2}{2}t}\phi(p'\mathrm{e}^{-t})$$

so that

$$S_t = -\frac{1}{2}\int_{p'} \phi'(p')\phi'(-p')\frac{{p'}^2}{K(p')}$$

becomes independent of t.

We now consider a generic theory:

$$\begin{aligned} S_t &= \sum_n \frac{1}{(2n)!}\int_{p_1,\cdots,p_{2n}} \phi(p_1)\cdots\phi(p_{2n})\, u_{2n}(t;p_1,\cdots,p_{2n}) \\ &\qquad \cdot(2\pi)^D\delta^{(D)}(p_1+\cdots+p_{2n}) \end{aligned}$$

Using ϕ' instead of ϕ, we obtain

$$\begin{aligned} S_t &= \sum_n \frac{1}{(2n)!}\int_{p_1,\cdots,p_{2n}} \mathrm{e}^{n(D+2)t}\phi'(p_1\mathrm{e}^t)\cdots\phi'(p_{2n}\mathrm{e}^t)\, u_{2n}(t;p_1,\cdots,p_{2n}) \\ &\qquad \cdot(2\pi)^D\delta^{(D)}(p_1+\cdots+p_{2n}) \end{aligned}$$

Hence, regarding S_t as a functional of ϕ', we obtain

$$\partial_t S_t = \int_{p'}\left\{p'_\mu\frac{\partial\phi'(p')}{\partial p'_\mu} + \frac{D+2}{2}\phi'(p')\right\}\frac{\delta S_t}{\delta\phi'(p')}$$

This is the modification due to rescaling.

We now note

$$\frac{\delta}{\delta\phi'(p')} = \mathrm{e}^{-\frac{D-2}{2}t}\frac{\delta}{\delta\phi(p'\mathrm{e}^{-t})}$$

HW#19: Derive this result. (Hint: Apply the above to $\phi'(q') = \mathrm{e}^{-\frac{D+2}{2}t}\phi(q'\mathrm{e}^{-t})$.)

Hence, the rescaling of space and field modifies the ERG differential equation to

$$\begin{aligned}\partial_t S_t &= \int_{p'} \left[\left\{ p'_\mu \frac{\partial \phi'(p')}{\partial p'_\mu} + \left(\frac{D+2}{2} + F'_t(p') \right) \phi'(p') \right\} \frac{\delta S_t}{\delta \phi'(p')} \right. \\ & \left. + G'_t(p') \frac{1}{2} \left\{ \frac{\delta S_t}{\delta \phi'(p')} \frac{\delta S_t}{\delta \phi'(-p')} + \frac{\delta^2 S_t}{\delta \phi'(p') \delta \phi'(-p')} \right\} \right]\end{aligned}$$

where

$$F'_t(p') \equiv F_t(p' \mathrm{e}^{-t}) = \frac{\Delta(p')}{K(p')} - \frac{\eta(t)}{2}$$

and

$$G'_t(p') \equiv \mathrm{e}^{-2t} G_t(p' \mathrm{e}^{-t})$$

Example 1 (Wilson)

$$G'_t(p) = 2 \left(\frac{\Delta(p)}{K(p)} + 1 - \frac{1}{2}\eta(t) \right)$$

Example 2 (Polchinski)

$$G'_t(p) = \frac{1}{p^2 + \mathrm{e}^{2t} m^2(t)} \left\{ \Delta(p) - \left(\eta(t) + \frac{b_m(t)}{p^2 + \mathrm{e}^{2t} m^2(t)} \right) K(p)(1 - K(p)) \right\}$$

where

$$\frac{d}{dt} \left(\mathrm{e}^{2t} m^2(t) \right) = (2 + b_m(t))\, \mathrm{e}^{2t} m^2(t)$$

From now, we call $\mathrm{e}^{2t} m^2(t)$ as $m^2(t)$. We will also omit the primes from ϕ' and F'_t, G'_t.

HW#20: Given

$$\begin{aligned}S_t[\phi] &= \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1,\cdots,p_{2n}} \phi(p_1) \cdots \phi(p_{2n})\, \mathcal{V}_{2n}(t; p_1, \cdots, p_{2n}) \\ & \times (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_{2n})\end{aligned}$$

before rescaling, derive

$$\begin{aligned}\underbrace{S'_t[\phi']}_{\equiv S_t[\phi]} &= \sum_{n=1}^{\infty} \frac{1}{(2n)!} \int_{p_1,\cdots,p_{2n}} \phi'(p_1) \cdots \phi'(p_{2n})\, \mathcal{V}'_{2n}(t; p_1, \cdots, p_{2n}) \\ & \times (2\pi)^D \delta^{(D)}(p_1 + \cdots + p_{2n})\end{aligned}$$

where

$$\mathcal{V}'_{2n}(t; p_1, \cdots, p_{2n}) = \mathrm{e}^{y_{2n} t} \mathcal{V}_{2n}(t; p_1 \mathrm{e}^{-t}, \cdots, p_{2n} \mathrm{e}^{-t})$$

and

$$y_{2n} \equiv D - n(D-2)$$

HW#21: Show that in the Polchinski formalism $\mathcal{V}_2$ and $\mathcal{V}_4$ satisfy the following ERG differential equations:

$$\begin{aligned}
\partial_t \mathcal{V}_2(t; p, -p) &= \eta(p^2+m^2) + b_m + \left(2 - \eta + 2u(p) - 2p^2 \frac{\partial}{\partial p^2}\right) \mathcal{V}_2(t; p, -p) \\
&\quad + \mathcal{V}_2 \frac{(\Delta - Ku)(p)}{p^2+m^2} \mathcal{V}_2 + \frac{1}{2} \int_q \frac{(\Delta - Ku)(q)}{q^2+m^2} \mathcal{V}_4(t; q, -q, p, -p) \\
\partial_t \mathcal{V}_4(t; p_1, \cdots, p_4) &= \left((4-D) - 2\eta + \sum_{i=1}^4 \left(u(p_i) - p_{i\mu} \frac{\partial}{\partial p_{i\mu}}\right)\right) \mathcal{V}_4 \\
&\quad + \mathcal{V}_4 \sum_{i=1}^4 \frac{(\Delta - Ku)(p_i)}{p_i^2+m^2} \mathcal{V}_2(p_i) + \frac{1}{2} \int_q \frac{(\Delta - Ku)(q)}{q^2+m^2} \mathcal{V}_6(t; q, -q, p_1, \cdots, p_4)
\end{aligned}$$

where

$$u(t; p) \equiv (1 - K(p)) \left(\eta(t) + \frac{b_m(t)}{p^2 + m^2(t)}\right)$$

5.5 The Wilson-Fisher fixed point for $D = 3$

For $D = 4$, the only fixed point of the ERG is the massless free theory. We will construct the fixed point explicitly for any D in the appendix at the end of this section. For $D = 3$, there is a non-trivial fixed point, discovered by Wilson and Fisher.

As the initial action at $t = 0$, let us consider

$$S_0 \equiv -\frac{1}{2} \int_p \phi(p)\phi(-p) \frac{p^2 + m_0^2}{K(p)} - \frac{\lambda_0}{4!} \int_{p_1,p_2,p_3} \phi(p_1)\phi(p_2)\phi(p_3)\phi(-p_1-p_2-p_3)$$

For a fixed $\lambda_0 > 0$, we consider starting the ERG trajectories with various values of m_0^2. For a generic choice of m_0^2, the theory is massive, and the ERG trajectory leads to an infinitely massive theory with no correlations since the physical mass increases exponentially along the ERG trajectory. However, for a specific value of $m_0^2 = m_{cr}^2(\lambda_0)$, the theory is massless, and the ERG trajectory goes into the Wilson-Fisher fixed point as $t \to \infty$. For $m_0^2 > m_{cr}^2(\lambda_0)$, the theory has $\mathbf{Z}_2$ invariance, but it is broken spontaneously for $m_0^2 < m_{cr}^2(\lambda_0)$.

In the modified Polchinski ERG, we have a running squared mass $m^2(t)$ whose initial value is given by $m^2(0) = m_0^2$. We expect that

$$m^2(t) \overset{t\to\infty}{\longrightarrow} m^{2*}$$

for $m_0^2 = m_{cr}^2(\lambda_0)$. Here m^{2*} is the squared mass parameter of the fixed point action.



Figure 12: At criticality $m_0^2 = m_{cr}^2(\lambda_0)$, the ERG trajectory leads to the Wilson-Fisher fixed point.

5.6 Perturbative determination of critical exponents

We finally discuss a perturbative construction of the Wilson-Fisher fixed point. Especially we wish to compute the anomalous dimensions (critical exponents) of the squared mass and ϕ using perturbation theory.

For perturbation theory, the modified Polchinski ERG differential equation is the most convenient.[18]

$$\begin{aligned}\partial_t S_t = & \int_p \left\{ p_\mu \frac{\partial \phi(p)}{\partial p_\mu} + \left(\frac{D+2}{2} - \frac{\eta(t)}{2} + \frac{\Delta(p)}{K(p)} \right) \phi(p) \right\} \frac{\delta S_t}{\delta \phi(p)} \\ & + \int_p \frac{1}{p^2+m^2(t)} \left\{ \Delta(p) - \left(\eta(t) + \frac{b_m(t)}{p^2+m^2(t)} \right) K(p)\,(1-K(p)) \right\} \\ & \qquad \times \frac{1}{2} \left(\frac{\delta S_t}{\delta \phi(p)} \frac{\delta S_t}{\delta \phi(-p)} + \frac{\delta^2 S_t}{\delta \phi(p) \delta \phi(-p)} \right)\end{aligned}$$

where $m^2(t)$ satisfies

$$\frac{d}{dt} m^2(t) = 2m^2(t) + b_m(t)$$

The two normalization conditions

$$\left\{ \begin{array}{rcl} \mathcal{V}_2(m^2; p, -p)\Big|_{p^2=0} & = & 0 \\ \frac{\partial}{\partial p^2} \mathcal{V}_2(m^2; p, -p)\Big|_{p^2=0} & = & 0 \end{array} \right.$$

determine $b_m(t)$ and $\eta(t)$ as follows:

$$\begin{aligned} & -b_m(t) - \eta(t) m^2(t) \\ & \quad = \frac{1}{2} \int_q \frac{1}{q^2+m^2(t)} \left\{ \Delta(q) - K(q)(1-K(q)) \left(\eta(t) + \frac{b_m(t)}{q^2+m^2(t)} \right) \right\} \end{aligned}$$

---

[18] Our scheme is not mass independent. There is a mass independent variation, and perhaps that is the simplest for perturbation theory.

$$\times\mathcal{V}_4(t;q,-q,0,0)$$

$$-\eta(t) = \frac{1}{2}\frac{\partial}{\partial p^2}\int_q \frac{1}{q^2+m^2(t)}\left\{\Delta(q) - K(q)(1-K(q))\left(\eta(t)+\frac{b_m(t)}{q^2+m^2}\right)\right\} \times\mathcal{V}_4(t;q,-q,p,-p)\Big|_{p^2=0}$$

Under the above ERG flow, the correlation functions obtain the following simple t-dependence:

$$\frac{1}{K(p\mathrm{e}^t)^2}\left(\langle\phi(p\mathrm{e}^t)\phi(-p\mathrm{e}^t)\rangle_{S(t)} - \frac{K(p\mathrm{e}^t)}{p^2\mathrm{e}^{2t}+m^2(t)}\right) + \frac{1}{p^2\mathrm{e}^{2t}+m^2(t)} = \exp\left[-2t+\int^t dt'\,\eta(t')\right]\underbrace{\langle\phi(p)\phi(-p)\rangle}_{t\text{-independent}}$$

$$\prod_{i=1}^{n>2}\frac{1}{K(p_i\mathrm{e}^t)}\cdot\langle\phi(p_1\mathrm{e}^t)\cdots\phi(p_n\mathrm{e}^t)\rangle_{S(t)} = \exp\left[t\left(D-n\frac{D+2}{2}\right)+\frac{n}{2}\int^t dt'\,\eta(t')\right]\cdot\underbrace{\langle\phi(p_1)\cdots\phi(p_n)\rangle}_{t\text{-independent}}$$

This is valid for any D, and we take $D=3$ here.

We wish to find the infrared (IR) fixed point using perturbation theory. It is convenient (and perhaps essential) to impose the UV renormalizability of the theory so that the theory can be characterized by only two parameters. If we solve the ERG differential equation using an initial condition such as given in the previous section, we need to keep track of every aspect of the action. It would be inconvenient for perturbative calculations.

The scalar theory in $D=3$ is a truly renormalizable theory, and UV renormalizability amounts to the asymptotic condition

$$S_t \overset{t\to-\infty}{\longrightarrow} S_G^* \equiv -\frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2}{K(p)}$$

where S_G^* is the gaussian fixed point theory, corresponding to the free massless theory.

The UV renormalizability familiar to us is formulated in terms of the action before rescaling of space: the six- and higher-point vertices should vanish asymptotically as $t\to-\infty$. Hence, using the result of HW#20, we obtain

$$\mathrm{e}^{(n-3)t}\mathcal{V}_{2n}(t;p_1\mathrm{e}^t,\cdots,p_{2n}\mathrm{e}^t) \overset{t\to-\infty}{\longrightarrow} 0$$

for $2n\geq 6$.

We can now parameterize the theory with $m^2(t)$ and the running coupling constant

$$\lambda(t)\equiv-\mathcal{V}_4(t;0,0,0,0)$$

These two parameters specify the UV asymptotic behavior of the action unambiguously, and hence the entire action is specified. Thus, we can write

$$S_t = S(m^2(t), \lambda(t))$$

The ERG differential equation gives

$$\frac{d\lambda(t)}{dt} = \lambda(t) + \beta(t)$$

where $\beta(t)$ is defined by

$$\begin{aligned} & -\beta(t) - 2\lambda(t) \cdot \eta(t) \\ & = \frac{1}{2} \int_q \frac{1}{q^2 + m^2(t)} \left\{ \Delta(q) - K(q)(1 - K(q)) \left(\eta(t) + \frac{b_m(t)}{q^2 + m^2(t)} \right) \right\} \\ & \qquad \times \mathcal{V}_6(t; q, -q, 0, 0, 0, 0) \end{aligned}$$

Since the action is completely characterized by the two parameters $m^2(t), \lambda(t)$, the ERG differential equation reduces to the two ordinary differential equations[19]:

$$\left\{ \begin{array}{rcl} \frac{dm^2}{dt} & = & 2m^2 + b_m(m^2, \lambda) \\ \frac{d\lambda}{dt} & = & \lambda + \beta(m^2, \lambda) \end{array} \right.$$

The fixed point (m^{2*}, λ^*) is determined by the two conditions:

$$\left\{ \begin{array}{lcl} 2m^{2*} + b_m(m^{2*}, \lambda*) & = & 0 \\ \lambda^* + \beta(m^{2*}, \lambda*) & = & 0 \end{array} \right.$$

To get the anomalous dimension of the squared mass, we linearize the RG equations:



Figure 13: ERG flows in the two-dimensional parameter space

$$\begin{array}{rcl} \frac{d}{dt}\left(m^2 - m^{2*}\right) & = & \left(2 + \frac{\partial b_m(m^2,\lambda)}{\partial m^2}\Big|^*\right)(m^2 - m^{2*}) + + \frac{\partial b_m(m^2,\lambda)}{\partial \lambda}\Big|^* (\lambda - \lambda^*) \\ \frac{d}{dt}\left(\lambda - \lambda^*\right) & = & \frac{\partial \beta(m^2,\lambda)}{\partial m^2}\Big|^* (m^2 - m^{2*}) + \frac{\partial \beta(m^2,\lambda)}{\partial \lambda}\Big|^* (\lambda - \lambda^*) \end{array}$$

---

[19] In a mass independent scheme, we will have $b_m(m^2, \lambda) = m^2 \beta_m(\lambda)$, and β depends only on λ.

This has two eigenvalues, one positive, and the other negative. The positive eigenvalue, $2+\beta_m^*$, gives the scale dimension of the relevant parameter (conjugate to something like ϕ^2), and the negative eigenvalue gives that of the least irrelevant parameter (conjugate to ϕ^4). We call β_m^* the **anomalous dimension** of the squared mass parameter.

Let us now calculate b_m, η, β, which are all functions of m^2, λ. At 1-loop, we obtain

$$\left\{\begin{array}{rcl} b_m(m^2,\lambda) &=& \frac{\lambda}{2}\int_q \frac{\Delta(q)}{q^2+m^2} \\ \beta(m^2,\lambda) &=& -3\lambda^2\int_q \frac{\Delta(q)(1-K(q))}{(q^2+m^2)^2} \\ \eta(m^2,\lambda) &=& 0 \end{array}\right.$$

Hence, the RG equations of the parameters become

$$\left\{\begin{array}{rcl} \frac{dm^2}{dt} &=& 2m^2 + \frac{\lambda}{2}\int_q \frac{\Delta(q)}{q^2+m^2} \\ \frac{d\lambda}{dt} &=& \lambda - 3\lambda^2\int_q \frac{\Delta(q)(1-K(q))}{(q^2+m^2)^2} \end{array}\right.$$

This implies

$$m^{2*} = \mathrm{O}(\lambda)$$

Hence, near the fixed point we can approximate the RG equations further as

$$\left\{\begin{array}{rcl} \frac{dm^2}{dt} &=& \left(2-\frac{\lambda}{2}\int_q \frac{\Delta(q)}{q^4}\right)m^2 + \frac{\lambda}{2}\int_q \frac{\Delta(q)}{q^2} \\ \frac{d\lambda}{dt} &=& \lambda - 3\lambda^2\int_q \frac{\Delta(q)(1-K(q))}{q^4} \end{array}\right.$$

Thus, we obtain

$$\left\{\begin{array}{rcl} \lambda^* &=& \frac{1}{3\int_q \frac{\Delta(q)(1-K(q))}{q^4}} \\ m^{2*} &=& -\frac{\lambda^*}{4}\int_q \frac{\Delta(q)}{q^2} \end{array}\right.$$

and the anomalous dimension of the squared mass is given by

$$\beta_m^* = -\frac{\lambda^*}{2}\int_q \frac{\Delta(q)}{q^4}$$

The fixed point values λ^*, m^{2*} have no physical meaning, and they depend on the choice of the cutoff function K. However, the anomalous dimension β_m^* is independent of the choice, and we can prove the independence by using an argument analogous to the one given in subsect. 2.12 for universality. Nevertheless, at any finite order, the perturbative result depends on K. To calculate β_m^*, we make the simplest choice

$$K(q) \to \theta(1-q^2)$$

Then, we obtain

$$\Delta(q) \equiv -2q^2\frac{d}{dq^2}K(q) = 2\delta(q^2-1)$$

Thus,

$$\int_q \frac{\Delta(q)}{q^4} = \frac{1}{(2\pi)^2}\int_0^\infty qdq^2\,\frac{1}{q^4}2\delta(q^2-1) = \frac{1}{2\pi^2}$$

Since

$$\Delta(q)K(q)^n = -2q^2 \frac{d}{dq^2} \frac{K(q)^{n+1}}{n+1}$$

we obtain

$$\int_q \frac{\Delta(q)K(q)^n}{q^4} = \frac{1}{n+1} \cdot \frac{1}{2\pi^2}$$

Thus, we obtain

$$\beta_m^* = -\lambda^* \int_q \frac{\Delta(q)(1 - K(q))}{q^4} = -\frac{1}{3}$$

This is the same as what we get from the ϵ expansions at the first order.

Appendix: The gaussian fixed point

In this appendix we would like to construct the gaussian fixed point explicitly using three formalisms: Polchinski with m^2, Polchinski with $m^2 = 0$, and Wilson. In all the formalisms, we look for a solution of the respective ERG differential equation corresponding to the free massive theory. The gaussian fixed point is the massless limit.

Polchinski with m^2

We look for a solution quadratic in fields:

$$S(t, m^2) = \frac{1}{2} \int_p \phi(p)\phi(-p) \left(-\frac{p^2 + m^2(t)}{K(p)} + \mathcal{V}_2(t; p, -p) \right)$$

In the absence of quartic interaction terms, both b_m and η vanish, and we obtain

$$m^2(t) = m^2 \mathrm{e}^{2t}$$

where m^2 is a constant. The vertex $\mathcal{V}_2$ satisfies the two conditions:

$$\begin{cases} \mathcal{V}_2(t, m^2; p, -p)\Big|_{p^2=0} &= 0 \\ \frac{\partial}{\partial p^2} \mathcal{V}_2(t, m^2; p, -p)\Big|_{p^2=0} &= 0 \end{cases}$$

and the ERG differential equation

$$\frac{\partial}{\partial t} \mathcal{V}_2(t, m^2; p, -p) = \left(2 - 2p^2 \frac{\partial}{\partial p^2} \right) \mathcal{V}_2(t, m^2; p, -p) + \frac{\Delta(p)}{p^2 + m^2 \mathrm{e}^{2t}} \mathcal{V}_2(t, m^2; p, -p)^2$$

The general solution is given by

$$\mathcal{V}_2(t, m^2; p, -p) = -\frac{\mathrm{e}^{2t}}{\frac{1}{p^4 \mathrm{e}^{-4t} f(p\mathrm{e}^{-t})} + \frac{1-K(p)}{p^2 \mathrm{e}^{-2t} + m^2}}$$

where $f(p)$ is an arbitrary function regular at $p^2 = 0$.

For any $m^2 \geq 0$, we obtain

$$\lim_{t \to \infty} \mathcal{V}_2(t; p, -p) = 0$$

so that the action is asymptotically given by the free action

$$S(t, m^2) \xrightarrow{t \gg 1} -\frac{1}{2} \int_p \phi(p)\phi(-p) \frac{p^2 + m^2 \mathrm{e}^{2t}}{K(p)}$$

The critical point is given by $m^2 = 0$, which gives the fixed point action:

$$\lim_{t \to \infty} S(t, 0) = S_G^* \equiv -\frac{1}{2} \int_p \phi(p)\phi(-p) \frac{p^2}{K(p)}$$

The continuum limit is obtained as

$$S_{\mathrm{free}}(\bar{m}^2) = \lim_{t \to \infty} S(t, m^2 = \bar{m}^2 \mathrm{e}^{-2t}) = -\frac{1}{2} \int_p \phi(p)\phi(-p) \frac{p^2 + \bar{m}^2}{K(p)}$$

Polchinski with $m^2 = 0$

We assume that the action is quadratic in fields:

$$S_t = \frac{1}{2}\int_p \phi(p)\phi(-p)\left(-\frac{p^2}{K(p)} + \mathcal{V}_2(t;p,-p)\right)$$

where the vertex $\mathcal{V}_2$ satisfies the condition

$$\frac{\partial}{\partial p^2}\mathcal{V}_2(t;p,-p)\Big|_{p^2=0} = 0$$

and the ERG differential equation

$$\frac{\partial}{\partial t}\mathcal{V}_2(t;p,-p) = \left(2 - 2p^2\frac{\partial}{\partial p^2}\right)\mathcal{V}_2(t;p,-p) + \frac{\Delta(p)}{p^2}\mathcal{V}_2(t;p,-p)^2$$

The general solution is given by

$$\mathcal{V}_2(t;p,-p) = -\frac{1}{\frac{1}{\mathrm{e}^{2t}(m^2+p^4\mathrm{e}^{-4t}f(p\mathrm{e}^{-t}))} + \frac{1-K(p)}{p^2}}$$

where $m^2 \geq 0$ is an arbitrary constant, and $f(p)$ is an arbitrary function of p^2 regular at $p^2 = 0$.

$m^2 = 0$ is the critical point:

$$\lim_{t\to\infty}\mathcal{V}_2(t;p,-p)\Big|_{m^2=0} = 0$$

so that the fixed point action is given by the same action as for the Polchinski with m^2:

$$S_G^* = -\frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2}{K(p)}$$

The massive free theory is obtained as the continuum limit:

$$\lim_{t\to\infty}\mathcal{V}_2(t;p,-p)\Big|_{m^2=\bar{m}^2\mathrm{e}^{-2t}} = -\frac{p^2\bar{m}^2}{p^2+\bar{m}^2(1-K(p))}$$

The action of the theory is given by

$$S_P(\bar{m}^2) \equiv -\frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2+\bar{m}^2}{K(p)}\frac{p^2}{p^2+\bar{m}^2(1-K(p))}$$

The cutoff-independent correlation function is obtained as

$$\frac{1}{p^2} + \frac{1}{K(p)^2}\left(\langle\phi(p)\phi(-p)\rangle_{S(\bar{m}^2)} - \frac{K(p)}{p^2}\right) = \frac{1}{p^2+\bar{m}^2}$$

Wilson

Let us assume a quadratic action:

$$S_t = \frac{1}{2} \int_p \phi(p)\phi(-p)\, u(t; p, -p)$$

In the Wilsonian case the anomalous dimension depends on both the quadratic and quartic terms. Hence, it may not vanish even for the quadratic theory. Hence, the ERG differential equation of Wilson gives

$$\begin{aligned} \partial_t u_2(t; p, -p) \quad &= \quad \left(2 - 2p^2 \frac{\partial}{\partial p^2} + 2\frac{\Delta(p)}{K(p)} - \eta(t)\right) u_2(t; p, -p) \\ &\quad + 2\left(\frac{\Delta(p)}{K(p)} + 1 - \frac{1}{2}\eta(t)\right) u_2(t; p, -p)^2 \end{aligned}$$

Given the initial condition

$$u_2(0; p, -p) = -\omega(p) = -\left(\omega(0) + p^2 + \cdots\right)$$

the solution is given by

$$u_2(t; p, -p) = -\frac{\mathrm{e}^{2t}\omega(p\mathrm{e}^{-t})}{\mathrm{e}^{2t}\omega(p\mathrm{e}^{-t}) + \mathrm{e}^{\int_0^t dt'\, \eta(t')} \frac{K(p)^2}{K(p\mathrm{e}^{-t})^2} \left(1 - \omega(p\mathrm{e}^{-t})\right)}$$

where

$$\mathrm{e}^{\int_0^t dt'\, \eta(t')} = \frac{1}{2\left(1 - \omega(0)\right)^2} \left\{1 - 2\mathrm{e}^{2t}\omega(0)\left(1 - \omega(0)\right) + \sqrt{1 - 4\mathrm{e}^{2t}\omega(0)\left(1 - \omega(0)\right)}\right\}$$

The solution is valid for any $t > 0$ if $\omega(0) = 0$. For $\omega(0) > 0$, however, the above solution is valid only for $t > 0$ that satisfies

$$4\mathrm{e}^{2t}\omega(0)\left(1 - \omega(0)\right) < 1$$

At the critical point $\omega(0) = 0$, we obtain

$$\lim_{t\to\infty} u_2(t; p, -p) = -\frac{p^2}{p^2 + K(p)^2}$$

corresponding to the fixed point action:

$$S_G^* = -\frac{1}{2} \int_p \phi(p)\phi(-p) \frac{p^2}{p^2 + K(p)^2}$$

Physically this is the same Gaussian fixed point as in the Polchinski with or without m^2, but the concrete form depends on the formalism. To obtain the massive continuum limit we choose the t-dependent initial condition

$$\omega(0) = \bar{m}^2 \mathrm{e}^{-2t}$$

so that

$$\lim_{t\to\infty} u_2(t;p,-p) = -\frac{p^2+\bar{m}^2}{p^2+\bar{m}^2+K(p)^2 z(\bar{m}^2)}$$

where

$$z(\bar{m}^2) \equiv \frac{1}{2}\left(1-2\bar{m}^2+\sqrt{1-4\bar{m}^2}\right)$$

Obviously, we must restrict $\bar{m}^2$ to

$$4\bar{m}^2 < 1$$

With the continuum limit action

$$S_W(\bar{m}^2) = -\frac{1}{2}\int_p \phi(p)\phi(-p)\frac{p^2+\bar{m}^2}{p^2+\bar{m}^2+K(p)^2 z(\bar{m}^2)}$$

the cutoff independent correlation function is obtained as

$$\frac{1}{K(p)^2}\left(\langle\phi(p)\phi(-p)\rangle_{S(\bar{m}^2)}-1\right) = \frac{z(\bar{m}^2)}{p^2+\bar{m}^2}$$

We get a mass dependent field normalization in this case.

The above continuum limit action S_W can be obtained from the continuum action S_P of the Polchinski formalism with $m^2=0$ as follows:

$$\begin{aligned} \mathrm{e}^{S_W[\phi]} \quad = \quad & \int [d\phi'] \exp\Bigg[S_P[\phi'] \\ & -\frac{1}{2}\int_p A(p)^2 \left(\phi(p)-\frac{B(p)}{A(p)}\phi'(p)\right)\left(\phi(-p)-\frac{B(p)}{A(p)}\phi'(-p)\right)\Bigg] \end{aligned}$$

where

$$\begin{cases} \frac{1}{A(p)^2} &= 1 - z(\bar{m}^2)\frac{K(p)(1-K(p))}{p^2} \\ \frac{B(p)^2}{A(p)^2} &= z(\bar{m}^2) \end{cases}$$

Acknowledgment

I would like to thank Drs. L. Akant and K. Ülker for giving me an opportunity to present the lectures in a stimulating environment. I also thank Bekir Can Lütfüoğlu for much help, and Dr. H. Gies for a list of review articles.

References

[1] Wilson K G, Kogut J 1974 *Phys. Rept.* **12** 75

[2] Polchinski J 1984 *Nucl. Phys.* **B231** 269

[3] Becchi C 1991 `arXiv:hep-th/9607188`

[4] Morris T R 1998 *Prog. Theor. Phys. Suppl.* **131** 395

[5] Litim D F, Pawlowski J M 1999 *The Exact Renormalization Group*, Eds. Krasnitz et al (World Scientific) 168

[6] Aoki K, 2000 *Int. J. Mod. Phys.* **B 14** 1249

[7] Bagnuls C, Bervillier C 2001 *Phys. Rept.* **348** 91

[8] Berges J, Tetradis N, Wetterich C 2002 *Phys. Rept.* **363** 223

[9] Polonyi J 2004 *Central Eur. J. Phys.* **1** 1

[10] Pawlowski J M 2005 `arXiv:hep-th/0512261`

[11] Gies H 2006 `arXiv:hep-ph/0611146`

[12] Delamotte B 2007 `arXiv:cond-mat/0702365`

[13] Morris T R 1994 *J. Phys.* **A9** 2411

[14] Sonoda H 2003 *Phys. Rev.* **D67** 065011

[15] Sonoda H 2007 *J. Phys.* **A40** 5733-5750

[16] Sonoda H 2007 *J. Phys.* **A40** 9675–9690

[17] Igarashi Y, Itoh K, Sonoda H 2007 *Prog. Theor. Phys.* **118** 121–134

[18] Higashi T, Itou E, Kugo T 2007 `arXiv:0709.1522`